arXiv:1409.5978v3 [math.DS] 5 Nov 2015

TYPICAL PROPERTIES OF PERIODIC TEICHMÜLLER GEODESICS

URSULA HAMENSTÄDT

Abstract. Consider a component $\mathcal{Q}$ of a stratum in the moduli space of area one abelian differentials on a surface of genus g. Call a property $\mathcal{P}$ for periodic orbits of the Teichmüller flow on $\mathcal{Q}$ *typical* if the growth rate of orbits with property $\mathcal{P}$ is maximal. Typical are: The logarithms of the eigenvalues of the symplectic matrix defined by the orbit are arbitrarily close to the Lyapunov exponents of $\mathcal{Q}$, and its trace field is a totally real splitting field of degree g over $\mathbb{Q}$. If $g \geq 3$ then periodic orbits whose $SL(2,\mathbb{R})$-orbit closure equals $\mathcal{Q}$ are typical. We show that for every $\ell \geq 2$, there are finitely many affine invariant manifolds in $\mathcal{Q}$ of rank ℓ which contain all affine invariant manifolds in $\mathcal{Q}$ of rank ℓ, and we establish finiteness of algebraically primitive Teichmüller curves.

1. Introduction

The mapping class group $\mathrm{Mod}(S)$ of a closed surface S of genus $g \geq 2$ acts by precomposition of marking on the Teichmüller space $\mathcal{T}(S)$ of marked complex structures on S. The action is properly discontinuous, with quotient the moduli space $\mathcal{M}_g$ of complex structures on S.

The goal of this paper is to describe properties of this action which are invariant under conjugation and hold true for conjugacy classes of mapping classes which are typical in the following sense.

The moduli space are area one abelian differentials on $\mathcal{S}$ decomposes into *strata* of differentials with zeros of given multiplicities. There is a natural $SL(2,\mathbb{R})$-action on any connected component $\mathcal{Q}$ of this moduli space. The action of the diagonal subgroup is the *Teichmüller flow* Φ^t.

Let Γ be the set of all periodic orbits for Φ^t on $\mathcal{Q}$. The length of a periodic orbit $\gamma \in \Gamma$ is denoted by $\ell(\gamma)$. Let $h > 0$ be the entropy of the unique Φ^t-invariant Borel probability measure on $\mathcal{Q}$ in the Lebesgue measure class [M82, V86]. As an application of [EMR12] (see also [EM11, H11]) we showed in [H13] that

$$\sharp\{\gamma \in \Gamma \mid \ell(\gamma) \leq R\} \sim \frac{e^{hR}}{hR}.$$

Date: November 5, 2015.

Keywords: Abelian differentials, Teichmüller flow, periodic orbits, Lyapunov exponents, trace fields, orbit closures, equidistribution

AMS subject classification: 37C40, 37C27, 30F60

Research partially supported by ERC grant 10160104.

Call a subset $\mathcal{A}$ of Γ *typical* if

$$\sharp\{\gamma \in \mathcal{A} \mid \ell(\gamma) \leq R\} \sim \frac{e^{hR}}{hR}.$$

Thus a subset of Γ is typical if its growth rate is maximal. The intersection of two typical subsets is typical.

A periodic orbit $\gamma \in \Gamma$ for Φ^t determines the conjugacy class of a pseudo-Anosov mapping class. Each mapping class acts on $H_1(S, \mathbb{Z})$. This defines a natural surjective [FM12] homomorphism

$$\Psi : \mathrm{Mod}(S) \to Sp(2g, \mathbb{Z}).$$

Thus a periodic orbit γ of Φ^t determines the conjugacy class $[A(\gamma)]$ of a matrix $A(\gamma) \in Sp(2g, \mathbb{Z})$.

Let

$$1 = \kappa_1 > \kappa_2 > \cdots > \kappa_g > 0$$

be the positive Lyapunov exponents of the Kontsevich Zorich cocycle with respect to the normalized Lebesgue measure on $\mathcal{Q}$. The fact that there are no multiplicities in the Lyapunov spectrum was shown in [AV07]. For $\gamma \in \Gamma$ let $\hat{\alpha}_i(\gamma)$ be the logarithm of the absolute value of the i-th eigenvalue of the matrix $A(\gamma)$ ordered in decreasing order and write $\alpha_i(\gamma) = \hat{\alpha}_i(\gamma)/\ell(\gamma)$. As $A(\gamma)$ is symplectic, with real leading eigenvalue $e^{\ell(\gamma)}$, we have

$$1 = \alpha_1(\gamma) \geq \cdots \geq \alpha_g(\gamma) \geq 0 \geq -\alpha_g(\gamma) \geq \cdots \geq -\alpha_1(\gamma) = -1.$$

As eigenvalues of matrices are invariant under conjugation, for $-g \leq i \leq g$ we obtain in this way a function $\alpha_i : \Gamma \to [-1, 1]$.

The characteristic polynomial of a symplectic matrix $A \in Sp(2g, \mathbb{Z})$ is a reciprocal polynomial of degree $2g$ with integral coefficients. Its roots define a number field K of degree at most $2g$ over $\mathbb{Q}$ which is a quadratic extension of the so-called *trace field* of A. The Galois group of K is isomorphic to a subgroup of the semi-direct product $(\mathbb{Z}/2\mathbb{Z})^g \rtimes \Sigma_g$ where Σ_g is the symmetric group in g variables (see [VV02] for details). The field K and the Galois group only depend on the conjugacy class of A.

For $\gamma \in \Gamma$ let $G(\gamma)$ be the Galois group of the number field defined by the conjugacy class $[A(\gamma)]$. We show

Theorem 1. (1) *For $\epsilon > 0$ the set $\{\gamma \in \Gamma \mid |\alpha_i(\gamma) - \kappa_i| < \epsilon\}$ $(1 \leq i \leq g)$ is typical.*

(2) *The set of all $\gamma \in \Gamma$ such that the trace field of $[A(\gamma)]$ is totally real, of degree g over $\mathbb{Q}$, and $G(\gamma) = (\mathbb{Z}/2\mathbb{Z})^g \rtimes \Sigma_g$ is typical.*

By the groundbreaking work of Eskin, Mirzakhani and Mohammadi [EMM13], the closure of an orbit for the $SL(2, \mathbb{R})$-action on $\mathcal{Q}$ is an affine invariant manifold. Examples of non-trivial orbit closures are arithmetic *Teichmüller curves*. They arise from branched covers of the torus, and they are dense in any stratum of abelian differentials. Other examples of orbit closures different from entire components of strata can be constructed using more general branched coverings.

Theorem 2. *Let $\mathcal{Q}$ be any component of a stratum in genus $g \geq 3$. Then the set of all $\gamma \in \Gamma$ whose $SL(2,\mathbb{R})$-orbit closure equals $\mathcal{Q}$ is typical.*

The *rank* of an affine invariant manifold $\mathcal{M}$ is defined by

$$\mathrm{rk}(\mathcal{M}) = \frac{1}{2}\dim(pT\mathcal{M})$$

where p is the projection of period coordinates into absolute cohomology [W15]. Teichmüller curves are affine invariant manifolds of rank one, and the rank of a component of a stratum equals g.

For $g = 2$, Theorem 2 is false in a very strong sense. Namely, McMullen [McM03] showed that in this case, the orbit closure of any periodic orbit is an affine invariant manifold of rank one. If the trace field K is quadratic then this orbit closure is mapped by the Torelli map into the Hilbert modular surface defined by K in the moduli space of principally polarized abelian varieties. Such a Hilbert modular surface is a quotient of $\mathbf{H}^2 \times \mathbf{H}^2$ by the lattice $PSL(2,\mathcal{O}_K)$ where $\mathcal{O}_K$ is the ring of algebraic integers in K. This is closely related to the classification of orbit closures in genus 2.

In higher genus, Apisa [Ap15] classified all orbit closures in hyperelliptic components of strata. For other components a classification of orbit closures is not available, but there is substantial recent progress towards a geometric understanding of orbit closures. We refer to the paper [LNW15] for an excellent recent overview of what is known and for references.

We contribute to these results by establishing a finiteness result for affine invariant submanifolds of rank at least two.

Theorem 3. *Let $g \geq 3$ and let $\mathcal{Q}$ be a component of a stratum in the moduli space of abelian differentials. For every $2 \leq \ell < g$, there are finitely many proper affine invariant submanifolds in $\mathcal{Q}$ of rank ℓ which contain every affine invariant submanifold of rank ℓ.*

Theorem 3 does *not* imply that there are only finitely many affine invariant manifolds of rank $2 \leq \ell < g$ in $\mathcal{Q}$. Namely, the result says nothing about affine invariant submanifolds of rank ℓ which are properly contained in another proper affine invariant manifold of the same rank. We conjecture that an affine invariant manifold $\mathcal{C}$ of any rank $\ell \geq 1$ contains only finitely many affine invariant manifolds of the same rank. The case when $\mathcal{C}$ is a component of a stratum (and hence the rank is maximal) is due to Mirkzakhani and Wright [MW15]. The paper [H15] covers the case when the dimension of $\mathcal{C}$ equals $2\mathrm{rk}(\mathcal{C}+1$. A partial result for rank one affine invariant manifolds is due to Lanneau, Nguyen and Wright [LNW15].

For a component $\mathcal{Q}$ of a stratum with a single zero, an affine invariant submanifold of rank ℓ does not contain a proper affine invariant manifold of the same rank and hence we obtain

Corollary. *A stratum of abelian differentials with a single zero only contains finitely many affine invariant manifolds of rank $\ell \geq 2$.*

In [H15], this result is extended to strata with two zeros.

Teichmüller curves are affine invariant manifolds of dimension 2. To each such Teichmüller curve, there is associated a *trace field* which is an algebraic number field of degree at most g over $\mathbb{Q}$. This trace field coincides with the trace field of every periodic orbit contained in the curve [KS00]. The Teichmüller curve is called *algebraically primitive* if the degree of its trace field equals g.

The stratum $\mathcal{H}(2)$ of abelian differentials with a single zero on a surface of genus 2 contains infinitely many algebraically primitive Teichmüller curves [McM03b]. Recently, Bainbridge, Habegger and Möller [BHM14] showed finiteness of algebraically primitive Teichmüller curves in any stratum in genus 3. Finiteness of algebraically primitive Teichmüller curves in strata of differentials with a single zero for surfaces of prime genus $g \geq 3$ was established in [MW15]. Our final result generalizes this to every stratum in every genus $g \geq 3$, with a different proof.

Theorem 4. *Any component $\mathcal{Q}$ of a stratum in genus $g \geq 3$ contains only finitely many algebraically primitive Teichmüller curves.*

Plan of the paper: In Section 2 we establish the first part of Theorem 1 as a fairly easy consequence of the results in [H13].

Section 3 introduces the idea of a monodromy group and its Zarisky closure and uses it to show that for every component $\mathcal{Q}$ of a stratum and every component $\tilde{\mathcal{Q}}$ of its lift to the Teichmüller space of marked abelian differentials, the image under the homomorphism Ψ of the stabilizer of $\tilde{\mathcal{Q}}$ in the mapping class group is Zariski dense in $Sp(2g, \mathbb{R})$. A local version of this result, group sieving and the first part of Theorem 1 then leads to the second part of Theorem 1.

In Section 4 we look at the local monodromy group of an affine invariant manifold and show that it is Zariski dense in the symplectic group of rank corresponding to the rank of the manifold. We then compare the Chern connection on the Hodge bundle to the Gauss Manin connection and prove Theorem 3. The proofs of Theorem 2 and Theorem 4 are contained in Section 5.

Acknowledgement: I am very grateful to Alex Wright for pointing out a mistake in an earlier version of this work. Thanks to Alex Eskin and Alex Wright for helpful discussions. Part of this work was carried out in spring 2015 while I participated in the program on higher Teichmüller theory at the MSRI in Berkeley. I gratefully acknowledge support by a grant from the NSF during my stay. When this work was written I learned that Eskin, Filip and Wright obtained a stronger version of Theorem 3.

2. Lyapunov exponents

In this section we consider any component $\mathcal{Q}$ of a stratum of area one abelian differentials. All the results in this section are equally valid for any affine invariant submanifold for the $SL(2, \mathbb{R})$-action [EMM13].

Denote by $\tilde{\mathcal{Q}}$ a component of the preimage of $\mathcal{Q}$ in the Teichmüller space of marked area one abelian differentials. Let $h > 0$ be the entropy of the Teichmüller flow Φ^t on $\mathcal{Q}$ with respect to the invariant Lebesgue measure λ.

Let $1 = \kappa_1 \geq \cdots \geq \kappa_g \geq 0$ be the nonnegative Lyapunov exponents of the Teichmüller flow on $\mathcal{Q}$. For components of strata, the exponents are all distinct and positive [AV07], but this need not be the case for more general affine invariant manifolds. We refer to [Au14] for a discussion and examples.

Let

$$\Gamma \subset \mathcal{Q}$$

be the countable collection of all periodic orbits for Φ^t contained in $\mathcal{Q}$. Denote by $\ell(\gamma)$ the period of $\gamma \in \Gamma$. The orbit γ determines a conjugacy class in $\mathrm{Mod}(S)$ of pseudo-Anosov elements. Let $\varphi \in \mathrm{Mod}(S)$ be such an element; then $A(\gamma) = \Psi(\varphi) \in \mathrm{Sp}(2g, \mathbb{Z})$ is determined by γ up to conjugation. In particular, the quotients by $\ell(\gamma)$ of the logarithms $1 = \alpha_1(\gamma) \geq \cdots \geq \alpha_g(\gamma) \geq 0$ of the g largest absolute values of the eigenvalues of the matrix $A(\gamma)$ ordered in decreasing order only depend on γ but not on any choices made.

Let $\epsilon > 0$. For $\gamma \in \Gamma$ define $\chi_\epsilon(\gamma) = 1$ if $|\alpha_i(\gamma) - \kappa_i| < \epsilon$ for every $i \in \{1, \ldots, g\}$ and define $\chi_\epsilon(\gamma) = 0$ otherwise.

For $R_1 < R_2$ let $\Gamma(R_1, R_2) \subset \Gamma$ be the set of all periodic orbits of prime period contained in the interval (R_1, R_2). For an open or closed subset V of $\mathcal{Q}$ define

$$H(V, R_1, R_2) = \sum_{\gamma \in \Gamma(R_1, R_2)} \int_\gamma \chi(V) \quad \text{and}$$

$$H_\epsilon(V, R_1, R_2) = \sum_{\gamma \in \Gamma(R_1, R_2)} \int_\gamma \chi(V)\chi_\epsilon(\gamma)$$

where $\chi(V)$ is the characteristic function of V. Clearly we have $H_\epsilon(V, R_1, R_2) \leq H(V, R_1, R_2)$ for all $\epsilon > 0$.

Call a point $q \in \mathcal{Q}$ *birecurrent* if it is contained in its own α- and ω limit set. By the Poincaré recurrence theorem, the set of birecurrent points in $\mathcal{Q}$ has full Lebesgue measure. In [H13] (Corollary 4.8 and Proposition 5.4) we showed

Proposition 2.1. *For every birecurrent point $q \in \mathcal{Q}$, for every neighborhood U of q in $\mathcal{Q}$ and for every $\delta > 0$ there is an open neighborhood $V \subset U$ of q in $\mathcal{Q}$ and a number $t_0 > 0$ such that*

$$H(V, R - t_0, R + t_0)e^{-hR} \in (2t_0\lambda(V)(1 - \delta), 2t_0\lambda(V)(1 + \delta))$$

for all sufficiently large $R > 0$.

Denote by $\mathcal{H} \to \mathcal{Q}$ the pull-back to $\mathcal{Q}$ of the flat bundle over moduli space whose fibre at x is the real first cohomology group $H^1(S, \mathbb{R})$. Thus the fibre of the bundle $\mathcal{H}$ naturally has the structure of a symplectic vector space of real dimension $2g$. The integral homology of S determines the *Gauss Manin connection* on $\mathcal{H}$. This connection is flat. Parallel transport for this connection defines a lift of the Teichmüller flow Φ^t to a flow Θ^t on $\mathcal{H}$ which preserves the symplectic structure. This flow extension is called the *Kontsevich Zorich cocycle* for Φ^t.

The first part of Theorem 1 is an immediate consequence of Proposition 2.1 and the following

Proposition 2.2. *For every birecurrent point $q \in \mathcal{Q}$, for every neighborhood U of q in $\mathcal{Q}$ and for every $\delta > 0$ there is an open neighborhood $V \subset U$ of q in $\mathcal{Q}$ and a number $t_0 > 0$ with the properties stated in Proposition 2.1 such that for every $\epsilon > 0$ we have*

$$\lim \inf_{R\to\infty} H_\epsilon(V, R - t_0, R + t_0)e^{-hR} \geq 2t_0\lambda(V)(1 - \delta).$$

Proof. Let $\| \|$ be the *Hodge norm* on the bundle $\mathcal{H} \to \mathcal{Q}$ (see [ABEM12] for definitions and the most important properties). For $z \in \mathcal{Q}$, for $1 \leq i \leq g$ and for $t > 0$ let

$$\zeta_i(t, z)$$

be the minimum of the operator norms of the restriction of $\Theta^t(z)$ to a symplectic subspace of the fibre $\mathcal{H}_z$ of real dimension $2(g - i + 1)$. Define

$$\kappa_i(t, z) = \frac{1}{t} \log \zeta_i(t, z).$$

Let $\delta > 0$. Since the Kontsevich Zorich cocycle is locally constant, Lemma 4.7 of [H13] together with Proposition 5.4 of [H13] imply that for every neighborhood U of q there is a nested sequence of neighborhoods $Z_0 \subset Z_1 \subset Z_2 \subset Z_3 \subset V \subset U$ of q, with Z_i closed and V open and contractible, and there is a number $R_0 > 0$ with the following properties.

Let $z \in Z_0$ and let $R > R_0$ be such that $\Phi^R z \in Z_1$. Let $\hat{E}$ be the connected component containing z of the intersection $\Phi^t V \cap V$.

(1) The length of the connected orbit segment of $\cup_{t\in\mathbb{R}}\Phi^t z \cap Z_1$ containing $\Phi^R z$ equals $2t_0$.
(2) $\lambda(Z_0) > (1 + \delta)^{-1}\lambda(V)$, and the Lebesgue measure of the intersection $\Phi^R Z_2 \cap Z_3 \cap \hat{E}$ is contained in the interval
$$[e^{-hR}\lambda(V)/(1 + \delta), e^{-hR}\lambda(V)(1 + \delta)].$$
(3) Connect $\Phi^R z$ to z by an arc in V and let η be the resulting loop. Then there is a unique periodic orbit γ for Φ^t of length at most $R + \delta$ which passes through Z_3 and intersects $\Phi^R Z_2 \cap Z_3 \cap \hat{E}$, and this orbit is contained in the δ-neighborhood of η for some fixed choice of a Riemannian metric on $\mathcal{Q}$.
(4)
$$|\kappa_i(R, z) - \alpha_i(\gamma)| \leq \epsilon/2.$$

By Oseledec's theorem and ergodicity, there is number $R(\epsilon) > R_0$ and a subset B of Z_0 of measure $\lambda(B) > \lambda(Z_0)(1 + \delta)^{-1}$ with the following property. Let $u \in B$ and let $R > R(\epsilon)$; then $|\kappa_i(R, u) - \kappa_i| \leq \epsilon/2$.

Since the Lebesgue measure is mixing for the Teichmüller flow, there is a number $R_1 > R(\epsilon)$ such that

$$\lambda(\Phi^R B \cap B) \geq \lambda(B)^2(1 + \delta)^{-1} \geq \lambda(Z_0)^2(1 + \delta)^{-3} \geq \lambda(V)^2(1 + \delta)^{-5}$$

for all $R \geq R_1$. By the estimate (2), this implies that the number of components of the intersection $\Phi^R V \cap V$ containing points in $\Phi^R B \cap B$ is at least $e^{hR}\lambda(V)(1+\delta)^{-6}$. By (3) above, for each such component there is a periodic orbit of Φ^t passing through $Z_3 \subset V$. The estimate (4) together with the definition of B yields that each such orbit γ satisfies $\chi_\epsilon(\gamma) = 1$. Thus each such component of intersection $\Phi^R V \cap V$ contributes $2t_0$ to the value $H_\epsilon(V, R - t_0, R + t_0)$. Together this shows the proposition. □

3. Galois groups

In this section we prove the second part of Theorem 1. Throughout the section, we denote by $\mathcal{Q}$ a component of a stratum in the moduli space of area one abelian differentials on S, and by $\tilde{\mathcal{Q}}$ a component of its preimage in the Teichmüller space of abelian differentials.

A periodic orbit of Φ^t on $\mathcal{Q}$ determines a conjugacy class in $Sp(2g, \mathbb{Z})$. However, it will be convenient to look at actual elements of $Sp(2g, \mathbb{Z})$ rather than at conjugacy classes.

Choose a birecurrent point $q \in \mathcal{Q}$. Let $\delta > 0$ and let $Z_0 \subset Z_1 \subset Z_2 \subset Z_3 \subset V \subset \mathcal{Q}$ be a nested family of neighborhoods of q in $\mathcal{Q}$ as constructed in [H13] (compare the proof of Proposition 2.2). All we need to know is that this family can be used to construct parametrized periodic orbits for Φ^t whose basepoints are contained in Z_3 as follows.

For some sufficiently large $R_0 > 0$ let $z \in Z_2$ and let $T > R_0$ be such that $\Phi^T z \in Z_2$. Connect $\Phi^T z$ to z by an arc entirely contained in V. Call the resulting closed curve a *characteristic curve*. This curve determines uniquely a periodic orbit γ of Φ^t which intersects Z_3 in an arc of length $2t_0$ and does not depend on choices made [H13]. Choose the midpoint of this intersection arc as a basepoint for γ and as an initial point for a parametrization of γ.

Let Γ_0 be the set of all parametrized periodic orbits of this form for points $z \in Z_0$ with $\Phi^T z \in Z_1$. By Lemma 4.7 of [H13], the map which associates to a component of $\Phi^T V \cap V$ containing points in $\Phi^T Z_0 \cap Z_1$ the corresponding parametrized periodic orbit in Γ_0 is a bijection.

The set V is contractible. Fix once and for all a lift $\tilde{V}$ of V to $\tilde{\mathcal{Q}}$. A periodic orbit γ which intersects Z_3 in an arc of length $2t_0$ lifts to a subarc of a flow line of the Teichmüller flow on $\tilde{\mathcal{Q}}$ with starting point in $\tilde{V}$. The endpoints of this arc are identified by a pseudo-Anosov element $\Omega(\gamma) \in \mathrm{Mod}(S)$.

The following was proved in [H13]. We refer to the proof of Proposition 2.2 for notations.

Proposition 3.1. *For $\gamma_1, \ldots, \gamma_k \in \Gamma_0$, there is a point $z \in Z_2$, and there are numbers $0 < t_1 < \cdots < t_k$ with the following properties.*

(1) $\Phi^{t_i} z \in Z_2$.

(2) *The characteristic curve ζ of the orbit segment $\{\Phi^t z \mid 0 \leq t \leq t_k\}$ is contained in the δ-neighborhood of a periodic orbit γ.*
(3) $\Omega(\gamma) = \Omega(\gamma_k) \circ \cdots \circ \Omega(\gamma_1)$.

Proof. In the case that the arcs γ_i are contained in a fixed compact invariant subset K for Φ^t and that the set V is chosen small in dependence of K, the lemma is identical with the slight weakening of the statement of Theorem 4.3 of [H10]. That the statement holds true in the form presented here is immediate from the construction of the set Z_3 in [H13]. □

As a consequence, the subsemigroup $\Omega(\Gamma_0)$ of $\mathrm{Mod}(S)$ generated by $\{\Omega(\gamma) \mid \gamma \in \Gamma_0\}$ consists of pseudo-Anosov elements whose corresponding periodic orbits pass through Z_2.

Our next goal is to obtain information on the image of this subsemigroup in $SL(2g, \mathbb{Z})$. For this we choose an odd prime p and study the image of $\Psi\Omega(\Gamma_0)$ under the natural reduction map

$$\Lambda_p : Sp(2g, \mathbb{Z}) \to Sp(2g, F_p)$$

where F_p is the field with p elements. Denote by ι the symplectic form on F_p^{2g} which is preserved by $Sp(2g, F_p)$.

The following lemma relies on the results in [Hl08]. For its formulation, define a *transvection* in $Sp(2g, F_p)$ to be a map $A \in Sp(2g, F_p)$ of order p which fixes a subspace of F_p^{2g} of codimension one and has determinant one (see [Hl08]). Any map of the form $\alpha \to \iota(\alpha, \beta)\beta$ for some $\beta \in F_p^{2g}$ where ι is the symplectic form is a transvection. We call this map a *transvection with β*.

Lemma 3.2. *Let $p \geq 3$ be an odd prime and let $G < Sp(2g, F_p)$ be a subgroup generated by $2g$ transvections with a set $\mathcal{E} = \{e_1, \dots, e_{2g}\} \subset F_p^{2g}$ which spans F_p^{2g}. Assume that there is no nontrivial partition $\mathcal{E} = \mathcal{E}_1 \cup \mathcal{E}_2$ so that $\iota(e_{i_1}, e_{i_2}) = 0$ for all $e_{i_j} \in \mathcal{E}_j$. Then $G = Sp(2g, F_p)$.*

Proof. For each i write $A_i(x) = x + \iota(x, e_i)e_i$. Let $G < Sp(2g, F_p)$ be the subgroup generated by the transvections $A_1, \dots, A_{2g}$. Since the vectors $e_1, \dots, e_{2g}$ span F_p^{2g}, the invariant subspaces of the transvections A_i intersect trivially.

We claim that the standard representation of G on F_p^{2g} is irreducible. Namely, assume to the contrary that there is an invariant subspace $W \subset F_p^{2g}$. Let $0 \neq w \in W$; then there is at least one i so that $\iota(w, e_i) \neq 0$. By invariance, we have $w + \iota(w, e_i)e_i \in W$ and hence $e_i \in W$ since F_p is a field.

As a consequence, W is spanned by some of the e_i, say by $e_{i_1}, \dots, e_{i_k}$, and if j is such that $\iota(e_{i_s}, e_j) \neq 0$ for some $s \leq k$ then $e_j \in W$. However, this implies that $W = F_p^{2g}$ by the assumption on the set $\mathcal{E} = \{e_i\}$.

To summarize, G is an irreducible subgroup of $Sp(2g, F_p)$ generated by transvections (where irreducible means that the standard representation of σ on F_p^{2g} is irreducible), and the order of each of these transvections is not divisible by 2. Theorem 3.1 of [Hl08] now yields that $G = Sp(2g, F_p)$ which is what we wanted to show. □

Remark 3.3. By Proposition 6.5 of [FM12], Lemma 3.2 is not true for $p = 2$.

A slightly weaker version of the following proposition which is sufficient for our purpose and holds true for all affine invariant manifolds will be established in Section 4. We include the version below because it is useful in other context and makes the strategy of the argument more transparent.

Proposition 3.4. (1) *For an odd prime $p \geq 3$ the subgroup of $Sp(2g, F_p)$ generated by $\{\Lambda_p \Psi(\Omega(\gamma)) \mid \gamma \in \Gamma_0\}$ equals the entire group $Sp(2g, F_p)$.*

(2) *If $\mathcal{Q}$ is a stratum of abelian differentials with at least one simple zero then the subgroup of $\mathrm{Mod}(S)$ generated by $\{\Omega(\gamma) \mid \gamma \in \Gamma_0\}$ equals the entire mapping class group.*

Proof. We begin with showing the second part of the proposition.

The mapping class group is generated by finitely many Dehn twists about simple closed curves. A specific generating system are the so-called *Humphries generators* (see p.112 of [FM12]). These generators are Dehn twists about the set of simple closed curves $a_1, \ldots, a_g, c_1, \ldots, c_{g-1}, m_1, m_2$ shown in the Figure 1.



Figure 1

Let $\mathcal{Q}$ be a stratum of abelian differentials with at least one simple zero. By [KZ03], $\mathcal{Q}$ is connected. For the second part of the proposition it suffices to show that each of the Humphries generators T_c is contained in the subgroup of $\mathrm{Mod}(S)$ generated by $\Omega(\gamma)$ ($\gamma \in \Gamma_0$). Here T_c denotes the positive Dehn twist about the simple closed curve c.

By [H11], to each component $\mathcal{Q}$ of a stratum there is associated a collection of *train tracks* which parametrize the component. If $\mathcal{U}$ is a component of a stratum and if $\mathcal{U}$ is contained in the closure of $\mathcal{Q}$ then any train track associated to $\mathcal{U}$ can be obtained from some train track associated to $\mathcal{Q}$ by removing some branches [H11].

Let now $\mathcal{U}$ be the stratum of differentials with one simple zero and one zero of order $2g-3$. We claim that there is a train track τ for $\mathcal{U}$ with the following property. For each of the Humphries generators T_c of $\mathrm{Mod}(S)$ and for $\sigma(c) \in \{1, -1\}$, we have $T_c^{\sigma(c)}(\tau) \prec \tau$ which means that the train track $T_c^{\sigma(c)}(\tau)$ is *carried* by τ.

Let as before ι be the intersection form on $H_1(S, \mathbb{Z})$. Denote by $[c]$ the homology class of an oriented simple closed curve c. Orient the curves in Figure 1 in such a way that for each i we have $\iota([c_{2i-1}], [a_{2i-1}]) = 1 = \iota([c_{2i-1}], [a_{2i}])$ and $\iota([c_{2i}], [a_{2i}]) = \iota([c_{2i}], [a_{2i+1}]) = 1$ and $\iota([a_2], [m_1]) = -1, \iota([a_3], [m_2]) = 1$. Construct from this oriented curve system a train track η by replacing each intersection of oriented curves by a large branch as shown in Figure 2. It is easy to check that η is orientable.



Figure 2

Each of the curves a_i, c_i, m_i is embedded in η, and the image of η under either the positive or the negative Dehn twist about any of these curves is carried by η. Namely, the curves are embedded as a subgraph which either consists of a single large branch and a single small branch (in the case of the curves a_1, a_g, m_1, m_2) or of two large branches and two small branches as shown in Figure 3 below, or of three large branches and three small branches (for the curves a_2, a_3). The collection



Figure 3

of these Dehn twists defines the system of Humphries generators for $\mathrm{Mod}(S)$.

The train track η has two complementary components, one of them is a four-gon. It is not hard to see that η is birecurrent and hence it is associated to the stratum $\mathcal{U}$ of differentials with one simple zero and one zero of order $2g-3$ [H11]. By construction, it has the properties required in the above claim.

A stratum $\mathcal{Q}$ of differentials with a simple zero contains $\mathcal{U}$ in its closure [KZ03]. There is a train track τ for $\mathcal{Q}$ so that η can be obtained from τ by removing some branches [H11], i.e. η is a subtrack of τ. We can choose τ in such a way that for each of the Humphries generators T_c, we have $T_c^{\sigma(c)}\tau \prec \tau$. Thus τ is a train track as required in the above claim.

Choose a pseudo-Anosov mapping class φ which admits τ as a train track expansion. This means that $\varphi\tau \prec \tau$. Assume that φ maps every branch of τ *onto* τ. This will guarantee that the periodic orbit γ defined by φ is contained in $\mathcal{Q}$ rather than in the boundary of $\mathcal{Q}$, see [H11]. Then for each of the Humphries generators $T_c^{\sigma(c)} \in \mathrm{Mod}(S)$, the composition $\varphi \circ T_c^{\sigma(c)}$ satisfies $(\varphi \circ T_c^{\sigma(c)})\tau \prec \tau$, moreover $\varphi \circ T_c^{\sigma(c)}$ maps every branch of τ *onto* τ. But this just means that $\varphi \circ T_c^{\sigma(c)}$ is a pseudo-Anosov mapping class which admits τ as a train track expansion. In particular, $\varphi \circ T_c^{\sigma(c)}$ determines a periodic orbit in $\mathcal{Q}$.

We next show that for every periodic orbit γ on $\mathcal{Q}$ defined by a pseudo-Anosov mapping class with train track expansion τ as above and for every neighborhood U of $\gamma(0)$ (see the proof of Proposition 2.2, the subgroup of $\mathrm{Mod}(S)$ generated by

the parametrized orbits in the set Γ_0 which is constructed from a nested set of neighborhoods $Z_0 \subset Z_2 \subset Z_2 \subset Z_3 \subset V \subset U$ of $\gamma(0)$ as in [H13] contains each of the Humphries generators $T_c^{\sigma(c)}$ and hence this group is just the entire mapping class group.

Namely, write $\beta = T_c^{\sigma(c)}$ for one of these Dehn twists. By the above discussion, for each k the mapping class $\varphi^k \circ \beta \circ \varphi^k$ is pseudo-Anosov, and it defines a periodic orbit $\gamma_{\varphi^k \circ \beta \circ \varphi^k}$ in $\mathcal{Q}$. As $k \to \infty$, the normalized Φ^t-invariant measures supported on the periodic orbits $\gamma_{\varphi^k \circ \beta \circ \varphi^k}$ converge weakly to the normalized Lebesgue measure on γ. Thus for sufficiently large k the periodic orbit $\gamma_{\varphi^k \circ \beta \circ \varphi^k}$ passes through Z_0 and hence it defines an element of Γ_0. Then $\beta = \varphi^{-k} \circ (\varphi^k \circ \beta \circ \varphi^k) \circ \varphi^{-k}$ is contained in the group generated by $\{\Omega(\gamma) \mid \gamma \in \Gamma_0\}$ as claimed. As a consequence, the second part of the proposition holds true for any choice of a neighborhood V of $\gamma(0)$.

To show the claim for an arbitrary open set U in $\mathcal{Q}$, observe that typical periodic orbits for Φ^t passing through U become equidistributed for the Lebesgue measure (see [H13] for a detailed discussion). In particular, they pass through Z_0 where the set Z_0 is chosen as above.

Now use the argument for the set Z_0 as follows. Let γ be a typical such orbit, parametrized that it belongs to the set Γ_0 where Γ_0 is defined using a nested sequence of subsets of U. Let $T > 0$ be such that $\Gamma(T) \in Z_0$. Let $\hat{\gamma}$ be the reparametrization of γ which satisfies $\hat{\gamma}(0) = \gamma(T)$. Apply the above construction to $\hat{\gamma}$ and observe that for a sufficiently large iterate of $\hat{\gamma}$ (and hence γ) the geodesic constructed as above passes through U and hence its reparametrization is contained in Γ_0. Thus the claim follows as before.

To show the first part, let c be an oriented non-separating simple closed curve on S which defines the homology class $[c]$. The action on homology of the positive Dehn twist T_c about c equals

$$T_c(\alpha) = \alpha + \iota(\alpha, [c])[c] \tag{1}$$

(Proposition 6.3 of [FM12]). In other words, a Dehn twist acts on $H_1(S, \mathbb{Z})$ as a transvection with $[c]$.

By [KZ03], for $g \geq 4$ the stratum $\mathcal{H}(2g-2)$ of differentials with a single zero consists of three connected components. One of these components is hyperelliptic, the other two components are distinguished by the parity of the spin structure they define. The stratum $\mathcal{H}(4)$ consists of two components, one of which is hyperelliptic, and $\mathcal{H}(2)$ is connected.

Any component $\mathcal{Q}$ of a stratum with more than one zero contains a component of $\mathcal{H}(2g-2)$ in its closure. Thus following the reasoning in the first part of this proof, it suffices to find for each of the components $\mathcal{V}$ of $\mathcal{H}(2g-2)$ a train track ζ associated to $\mathcal{V}$ with the following property. Let $p \geq 3$ be an odd prime. Then the subgroup of $Sp(2g, F_p)$ which is generated by those transvections in $Sp(2g, F_p)$ which are images under $\Lambda_p \circ \Psi$ of Dehn twists about embedded curves c in ζ with $T_c^{\sigma(c)}\zeta \prec \zeta$ ($\sigma \in \{-1, 1\}$) is all of $Sp(2g, F_p)$.

To this end let again η be the train track constructed in the beginning of this proof from the Humphries generators of $\mathrm{Mod}(S)$. Let ζ be the train track obtained from η by removing the small branch contained in the curve m_2. This train track is orientable, filling, birecurrent and if $g \geq 3$, then it is not invariant under a hyperelliptic involution. Hence it corresponds to one of the two non-hyperelliptic components of $\mathcal{H}(2g-2)$. For each curve $c \in \{c_i, a_i, m_1\}$ we have $T_c^{\sigma(c)}\zeta \prec \zeta$. Now for any choice of orientations of the curves in $\{c_i, a_i, m_1\}$, the homology classes $\{[c_i], [a_i], [m_1]\}$ are a basis of $H_1(S, \mathbb{Z})$. Moreover, any two curves intersect in at most one point and their union is a connected subset of S. This implies that the transvections by the homology classes of these curves satisfy the assumptions in Lemma 3.2. As a consequence, for each odd prime $p \geq 3$ these transvections generate $Sp(2g, F_p)$. This is what we wanted to show.

The other two components of $\mathcal{H}(2g-2)$ are treated in the same way. A train track for the hyperelliptic component of $\mathcal{H}(2g-2)$ can be constructed from the following curve system. Remove the curves m_1, m_2 from the Humphries generators and add a simple closed curve which intersects a_1 in a single point and does not intersect any other of the curves shown in Figure 1. The resulting set of curves is clearly invariant under the hyperelliptic involution. The same argument as for the non-hyperelliptic components completes the proof of the proposition.

A train track for the second non-hyperelliptic component can be obtained from the following curve system. Remove from the curve system for the hyperelliptic component the curve c_{g-1} and replace it by a curve which intersects both a_{g-2} and a_g in a single point and does not intersect any other curve. □

A more general version of the first part of the following corollary is needed in Section 4, the second part is of independent interest. For its formulation, call a component $\mathcal{Q}$ of a stratum *locally Zarisky dense* if the following holds true. Let U be any open subset of $\mathcal{Q}$ and let $\Omega(\Gamma_0)$ be the subsemigroup of $\mathrm{Mod}(S)$ generated in the sense discussed above by the periodic orbits for Φ^t passing through U. We require that the subgroup of $Sp(2g, \mathbb{Z})$ generated by $\Psi\Omega(\Gamma_0)$ is Zariski dense in $Sp(2g, \mathbb{R})$.

Corollary 3.5. (1) *A component $\mathcal{Q}$ of a stratum is locally Zariski dense.*
(2) *If $\mathcal{Q}$ is a stratum of differentials with a simple zero then the preimage of $\mathcal{Q}$ in the Teichmüller space of area one abelian differentials is connected.*

Proof. The first part of the corollary follows from the first part of Proposition 3.4 and the well known fact that a subgroup of $Sp(2g, \mathbb{Z})$ which surjects onto $Sp(2g, F_p)$ for all odd primes $p \geq 3$ is Zariski dense in $Sp(2g, \mathbb{R})$ (see [Lu99] for more and for references).

Now let $\mathcal{Q}$ be a stratum with a simple zero and let $\tilde{\mathcal{Q}}_0$ be a component of $\tilde{\mathcal{Q}}$. By the second part of Proposition 3.4, the stabilizer of $\tilde{\mathcal{Q}}_0$ in the mapping class group equals the entire mapping class group. As the components of the preimage of $\mathcal{Q}$ are permuted by the mapping class group, the second part of the corollary follows. □

Remark 3.6. More generally, one can ask about the orbifold fundamental group of the closure of a component $\mathcal{Q}$ of a stratum, or, equivalently, about the stabilizer in $\mathrm{Mod}(S)$ of a component of the preimage of $\mathcal{Q}$ in the moduli space of abelian differentials. We conjecture that for non-hyperelliptic components with at least two zeros, this stabilizer has finite index in $\mathrm{Mod}(S)$. We also expect that for components of strata with a single zero, the stabilizer has infinite index in $\mathrm{Mod}(S)$. For many components, we do not even know the image of the stabilizer in the symplectic group. For the hyperelliptic component, this image is a proper subgroup of $Sp(2g,\mathbb{Z})$.

For a periodic orbit γ for Φ^t let δ_γ be the Φ^t-invariant measure supported on γ whose total mass equals the period $\ell(\gamma)$ of γ. Recall that periodic orbits in the set Γ_0 are parametrized, so a single unparametrized periodic orbit may give rise to many different elements of Γ_0. The next proposition is the main technical tool towards the proof of the second part of Theorem 1. For its formulation, let $N(p)$ be the number of elements of $Sp(2g,F_p)$.

Proposition 3.7. *Let $B \in Sp(2g,F_p)$ be arbitrary and define*

$$\mathcal{B}(R,B) = \{\gamma \in \Gamma_0 \mid \ell(\gamma) \leq R, \Lambda_p \circ \Psi \circ \Omega(\gamma) = B\}.$$

Then as $R \to \infty$,

$$\sharp\mathcal{B}(R,B) \sim \frac{e^{hR}\lambda(Z_0)}{2ht_0N(p)}$$

independent of B up to a multiplicative error which only depends on Z_0 and which can be chosen arbitrarily close to one.

Proof. We show first that there is a number $a > 0$ such that

$$\sharp\mathcal{B}(R,B) \geq ae^{hR}$$

for all $B \in Sp(2g,F_p)$ and for all sufficiently large R.

Let $R_0 > 0, \delta > 0$ and let $Z_0 \subset Z_1 \subset Z_2 \subset Z_3 \subset V$ be sets as in the beginning of this section. Let $\mathcal{G}$ be the subsemigroup of $\mathrm{Mod}(S)$ generated by $\{\Omega(\gamma) \mid \gamma \in \Gamma_0\}$. By Lemma 3.1, this semigroup consists of pseudo-Anosov elements, and each $\rho \in \mathcal{G}$ is represented by a parametrized periodic orbit ζ for Φ^t which intersects Z_3 in a segment of length $2t_0$ containing $\zeta(0)$ as its midpoint. Vice versa, every periodic orbit which passes through Z_0 admits a parametrization so that the corresponding element of $\mathrm{Mod}(S)$ is contained in $\mathcal{G}$. Then $\Psi(\mathcal{G})$ is a subsemigroup of $Sp(2g,\mathbb{Z})$. This semigroup is mapped by Λ_p to a subsemigroup G_p of the finite group $Sp(2g,F_p)$.

As $Sp(2g,F_p)$ is finite, for every $A \in Sp(2g,F_p)$ there is some $\ell \geq 1$ such that $A^\ell = A^{-1}$. As a consequence, for all $x, y \in G_p$ we have $xy^{-1} \in G_p$ as well and hence $G_p < Sp(2g,F_p)$ is a group. Thus G_p equals the group generated by $\{\Lambda_p\Psi\Omega(\gamma) \mid \gamma \in \Gamma_0\}$. By the first part of Lemma 3.4, we have $G_p = Sp(2g,F_p)$.

Since $Sp(2g,F_p)$ is a *finite* group and the above argument applies to every open subset of $\mathcal{Q}$, there is a number $\hat{R} > R_0$ with the following property. Let $A \in Sp(2g,F_p)$ be arbitrary. Then there is some $z \in Z_0$ with $\Phi^t z \in Z_1$ for some $t \in (R_0, \hat{R})$ which determines a parametrized periodic orbit $\gamma(A) \in \Gamma_0$ with $\Lambda_p\Psi(\Omega(\gamma(A))) = A$.

Write $Z = Z_0$. Let $v \in Z$ be such that $\Phi^T v \in Z, \Phi^{T+U} v \in Z$ for some $T, U > R_0$. By Lemma 3.1, the pseudo-orbits $\{\Phi^t v \mid 0 \leq t \leq T\}$ and $\{\Phi^t v \mid T \leq t \leq T+U\}$ determine two parametrized periodic orbits γ_1, γ_2 for Φ^t which define elements $\Lambda_p\Psi(\Omega(\gamma_1)), \Lambda_p\Psi(\Omega(\gamma_2)) \in Sp(2g, F_p)$. Let $\gamma = \gamma_2 \hat{\circ} \gamma_1$ be the periodic orbit for $\{\Phi^t v \mid 0 \leq t \leq T+U\}$. The notation $\gamma = \gamma_2 \hat{\circ} \gamma_1$ indicates that the element γ of $\mathrm{Mod}(S)$ defined by γ is the product of the elements of $\mathrm{Mod}(S)$ defined by γ_1 and γ_2. We have

$$\Lambda_p\Psi(\Omega(\gamma_2 \hat{\circ} \gamma_1)) = \Lambda_p\Psi(\Omega(\gamma_2)) \circ \Lambda_p\Psi(\Omega(\gamma_1)).$$

Fix an element $B \in Sp(2g, F_p)$. If $A \in Sp(2g, F_p)$ is arbitrary and if $\zeta \in \Gamma_0$ is such that $\Lambda_p\Psi(\Omega(\zeta)) = A$ then $\Lambda_p\Psi(\Omega(\gamma(BA^{-1}) \hat{\circ} \zeta)) = B$. In particular, by Proposition 2.1, for sufficiently large $R > R_0$ the number of parametrized periodic orbits $\gamma \in \Gamma_0$ with $\ell(\gamma) \leq R + \hat{R}$ and $\Lambda_p\Psi(\Omega(\gamma)) = B$ is not smaller than the number of orbits in Γ_0 of length at most R. For large enough R, this number can be estimated as follows.

Using Proposition 2.1, for large R the volume of Z with respect to the sum of the measures supported on periodic orbits of Φ^t of length in the interval $[R - t_0, R + t_0]$ is contained in the interval

$$[2t_0 e^{hR} \lambda(Z)(1-\delta), 2t_0 e^{hR} \lambda(Z)(1+\delta)].$$

A periodic orbit intersects the set Z in arcs of length $2t_0$ [H13]. As each such intersection component defines an element of Γ_0 (recall that Γ_0 consists of parametrized orbits), we conclude that for large enough R the number of elements of Γ_0 of length contained in $[R - t_0, R + t_0]$ roughly equals $e^{hR}\lambda(Z)$. Summation yields that the number of elements of Γ_0 of length at most $2kt_0$ roughly equals

$$\sum_{i=1}^{k} e^{h(2i-1)t_0} \lambda(Z) \sim e^{2hkt_0} \lambda(Z)/2ht_0.$$

As a consequence, up to passing to a subsequence, there is a number $a > 0$ not depending on B such that the measures

$$he^{-hR} \sum_{\gamma \in \mathcal{B}(R,B)} \delta_\gamma \chi[\gamma(-t_0), \gamma(t_0)]$$

converge to a measure λ_B on Z of toal mass roughly contained in $[a\lambda(Z), \lambda(Z)]$ (there is a multiplicative error depending on the a priori chosen constant $\delta > 0$ which tends to one as $\delta \to 0$). Here as before, δ_γ is the Φ^t-invariant measure of total mass $\ell(\gamma)$ on γ. Since periodic orbits are equidistributed for the Lebesgue measure [H13] and every periodic orbit of length R can be concatenated with one of finitely many orbits to yield an orbit in $\mathcal{B}(R, B)$, this measure is a multiple of the Lebesgue measure (see [H13] for details).

By construction, we have

$$\lambda_B(Z) = c(B)\lambda(Z) \tag{2}$$

for a number $c(B) \in [a, 1]$ and such that $\sum_B c(B) = 1$.

Our goal is to show that $c(B)$ is independent of B. To this end recall that the Lebesgue measure λ is mixing of all orders [M82]. In particular, for large enough numbers $R, S > 0$ we have

$$\lambda(Z \cap \Phi^R Z \cap \Phi^{R+S}(Z)) \sim \lambda(Z)^3$$

and therefore $\lambda_B(Z \cap \Phi^R Z \cap \Phi^{R+S} Z) \sim c(B)\lambda(Z)^3$.

For $R > 0, T > 0$ and $B \in Sp(2g, F_p)$ let $\Gamma(R, T, B, Z)$ be the set of all parametrized periodic orbits $\gamma \in \Gamma_0$ for Φ^t with the following properties.

(1) The length of γ is contained in the interval $[R + T - t_0, R + T + t_0]$.
(2) γ is determined by some $v \in Z$, and there is a number $U \in [R - t_0, R + t_0]$ such that $\Phi^U v \in Z$.
(3) $\Lambda_p \Psi(\Omega(\gamma)) = B$.

Define similarly a set $\Gamma(R + T, B, Z)$ where we only require properties (1) and (3) above. It follows from the above discussion (compare [H13]) that

$$\sharp\Gamma(R + T, B, Z) \sim c(B)\lambda(Z)e^{h(R+T)}/2ht_0$$

for large enough R and similarly

$$\sharp\Gamma(R, T, B, Z) \sim c(B)\lambda(Z)^2 e^{h(R+T)}/(2ht_0)^2.$$

Each orbit $\gamma \in \Gamma(R, T, B, Z)$ can be represented in the form $\gamma = \gamma_2 \hat{\circ} \gamma_1$ for some $\gamma_1 \in \Gamma(R, A, Z)$ and some $\gamma_2 \in \Gamma(T, BA^{-1}, Z)$ so that $CA = B$. Moreover, if $\gamma_1 \in \Gamma(R, A, Z)$ for some $A \in Sp(2g, F_p)$ then for any $\gamma_2 \in \Gamma(T, BA^{-1}, Z)$ we have $\gamma_2 \hat{\circ} \gamma_1 \in \Gamma(R, T, B, Z)$.

As a consequence, for an arbitrarily chosen $\epsilon > 0$ and for sufficiently large $R > 0$, as $T \to \infty$ we observe that

$$\begin{aligned}\sharp\Gamma(R, T, B, Z) &= \sum_{A \in Sp(2g, F_p)} \sharp\Gamma(T, BA^{-1}, Z)\sharp\Gamma(R, A, Z) \\ &\sim \frac{1}{(2ht_0)^2}\lambda(Z)^2 e^{h(R+T)} \sum_{A \in Sp(2g, F_p)} c(BA^{-1})c(A).\end{aligned}$$

Now let $B \in Sp(2g, F_p)$ be such that $c(B) = \min\{c(A) \mid A\}$. Such an element exists since $Sp(2g, F_p)$ is finite. Since $\sum_A c(BA^{-1}) = 1$ we have $\sharp\Gamma(R, T, B, Z) \sim c(B)\lambda(Z)^2 e^{h(R+T)}/(2ht_0)^2$ only if $c(A) = \frac{1}{N(p)}$ for all A. The proposition follows. □

Remark 3.8. Proposition 3.7 can be viewed as an equidistribution result for conjugacy classes elements in $Sp(2g, F_p)$ defined by periodic orbits of the Teichmüller flow. The main difficulty lies in the fact that periodic orbits represent conjugacy classes in the mapping class group rather than actual elements, moreover we look at periodic orbits in strata rather than at all periodic orbits. Mixing of the Teichmüller flow replaces the familiar equidistribution of random walks on finite connected graphs.

Now we are ready to complete the proof of the second part of Theorem 1. To this end recall that the characteristic polynomial of a symplectic matrix $A \in Sp(2g, \mathbb{Z})$ is reciprocal of degree $2g$. The roots of such a polynomial come in pairs. The Galois group of the number field defined by the polynomial is a subgroup of the semidirect product

$$(\mathbb{Z}/2\mathbb{Z})^g \rtimes S_g$$

where S_g is the symmetric group in g elements (see [VV02] for a nice account on this classical fact), and S_g acts on $(\mathbb{Z}/2\mathbb{Z})^g$ by permutation of the factors.

In the sequel we call the Galois group of the field defined by the characteristic polynomial of a matrix $A \in Sp(2g, \mathbb{Z})$ simply the Galois group of A. It only depends on the conjugacy class of A. We say that the Galois group of A is *full* if it coincides with $(\mathbb{Z}/2\mathbb{Z})^g \rtimes S_g$.

Having a full Galois group makes also sense for an element in $Sp(2g, F_p)$. We use this as in [R08] as follows.

Let again $p \geq 5$ be a prime and let $N(p)$ be the number of elements of $Sp(2g, F_p)$. By Proposition 3.7, for large enough R and every $B \in Sp(2g, F_p)$, the number of orbits $\gamma \in \Gamma_0$ of length at most R with $\Lambda_p \circ \Psi \circ \Omega(\gamma) = B$ roughly equals $\frac{e^{hR}}{N(p)} \lambda(Z_0)/2ht_0$. On the other hand, if we denote by $R_p(2g)$ the subset of $Sp(2g, F_p)$ of elements with reducible characteristic polynomial then

$$\frac{|R_p(2g)|}{N(p)} < 1 - \frac{1}{3g}$$

(see Theorem 6.2 of [R08] for a reference to this classical result of Borel).

Now we follow the proof of Theorem 6.2 of [R08]. Namely, let $p_1, \dots, p_k$ be k distinct primes, and let $K = p_1 \cdots p_k$. Then the reduction $\Lambda_K(A)$ modulo K of any element $A \in Sp(2g, \mathbb{Z})$ is defined, and we have

$$\Lambda_K(A) = \Lambda_{p_1}(A) \times \cdots \times \Lambda_{p_k}(A).$$

By Proposition 3.7 (or, rather, Proposition 3.7 and its proof which is valid for any finite quotient of $Sp(2g, \mathbb{Z})$ onto which the group generated by $\Psi\Omega(\Gamma_0)$ surjects), for large enough R, for a periodic orbit γ of length at most R the probability that the Galois group of the characteristic polynomial of $A(\gamma)$ is *not* full is at most of the order of $(1 - \frac{1}{3g})^k$. Now as $k \to \infty$, we conclude that the Galois group of a typical periodic orbit for Φ^t is full.

Let $\omega \in \tilde{\mathcal{Q}}$ be a lift of a point on a typical periodic orbit γ for Φ^t. The periods of ω define an abelian subgroup $\Lambda = \omega(H_1(S, \mathbb{Z}))$ of $\mathbb{C}$ of rank two. Let $e_1, e_2 \in \Lambda$ be two points which are linearly independent over $\mathbb{R}$. Let K be the smallest subfield of $\mathbb{R}$ such that every element of Λ can be written as $ae_1 + be_2$, with $a, b \in K$; then $\Lambda \otimes_K K = K^2$. If we write $T = \Psi(A(\gamma)) + \Psi(A(\gamma))^{-1}$, then the field K also is the field of the characteristic polynomial of T. We call K the *trace field* of γ (see the appendix of [KS00] for more details).

Definition 3.9. The periodic orbit γ is called *algebraically primitive* if the trace field K of γ is a totally real number field of degree g over $\mathbb{Q}$, with maximal Galois group.

The following corollary completes the proof of the second part of Theorem 1.

Corollary 3.10. *Algebraically primitive periodic orbits for Φ^t are typical.*

Proof. Since the Lyapunov spectrum of $\mathcal{Q}$ is simple [AV07], the first part of Theorem 1 implies that for a typical periodic orbit γ the absolute values of the eigenvalues of $A(\gamma)$ are pairwise distinct and hence all eigenvalues are real. Thus by the discussion following Proposition 3.7, we only have to show that for a symplectic matrix $A \in Sp(2g, \mathbb{R})$ with $2g$ distinct real eigenvalues r_i, r_i^{-1} $(i \leq g, r_i > 1)$ the field defined by $A + A^{-1}$ is totally real. However, this is immediate from the fact that the roots of the polynomial defining the trace field are of the form $r_i + r_i^{-1}$ where r_i are the roots of the characteristic polynomial of A. □

4. Affine invariant manifolds

Throughout this section we assume that $g \geq 3$.

A point in *Siegel upper half-space* $\mathcal{D}_g = \mathrm{Sp}(2g, \mathbb{R})/U(g)$ is a principally polarized abelian variety of dimension g. Here as usual, $U(g)$ denotes the unitary group of rank g. There is a natural rank g holomorphic vector bundle $\tilde{\mathcal{V}} \to \mathcal{D}_g$ whose fibre over y is just the complex vector space defining y. The polarization and the complex structure define a Hermitean metric h on $\tilde{\mathcal{V}}$. The group $Sp(2g, \mathbb{R})$ acts from the left on the bundle $\tilde{\mathcal{V}}$ as a group of bundle automorphisms preserving the polarization and the complex structure, and hence this action preserves the Hermitean metric. Thus the bundle $\tilde{\mathcal{V}}$ projects to a holomorphic Hermitean (orbifold) vector bundle

$$\mathcal{V} \to Sp(2g, \mathbb{Z}) \backslash Sp(2g, \mathbb{R})/U(g) = \mathcal{A}_g.$$

The Hermitean holomorphic vector bundle $\mathcal{V}$ admits a unique *Chern connection* ∇ (see e.g. [GH78]). The Chern connection defines parallel transport of the fibres of $\mathcal{V}$ along smooth curves in $\mathcal{A}_g$. As the Chern connection is Hermitean, parallel transport preserves the metric.

Let again $\mathcal{M}_g$ be the moduli space of closed Riemann surfaces of genus g. The *Hodge bundle*

$$\Pi : \mathcal{H} \to \mathcal{M}_g$$

is the pullback of the holomorphic bundle $\mathcal{V} \to \mathcal{A}_g$ under the *Torelli map*

$$\mathcal{J} : \mathcal{M}_g \to \mathcal{A}_g = Sp(2g, \mathbb{Z}) \backslash Sp(2g, \mathbb{R})/U(g).$$

As the Torelli map is holomorphic, $\mathcal{H}$ is a g-dimensional holomorphic vector bundle on $\mathcal{M}_g$ (in the orbifold sense). Its fibre over $x \in \mathcal{M}_g$ can be identified with the vector space of holomorphic one-forms on x. The natural Hermitean inner product on $\mathcal{H}$ is given by

$$(\omega, \zeta) = \frac{i}{2} \int \omega \wedge \overline{\zeta}.$$

With this interpretation, the sphere bundle in $\mathcal{H}$ for the inner product $(,)$ is just the moduli space of area one abelian differentials.

The bundle $\mathcal{H}$ admits a natural flat connection. First, each point $x \in \mathcal{M}_g$ determines a complex structure J_x on $H^1(S, \mathbb{R})$. Namely, each cohomology class

$\alpha \in H^1(S,\mathbb{R})$ can be represented by a unique harmonic one-form for the complex structure x, and this one-form is the real part of a unique holomorphic one-form ω on x. The imaginary part of this differential is a harmonic one-form which represents the cohomology class $J_x\alpha$. The complex structure J_x is *compatible* with the intersection form ι, i.e. we have $\iota(J_x\alpha, J_x\beta) = \iota(\alpha,\beta)$ for all $\alpha,\beta \in H^1(S,\mathbb{R})$.

In this way the (real) vector space $H^1(S,\mathbb{R})$ equipped with the complex structure J_x can naturally be identified with the fibre $\mathcal{H}_x$ of the Hodge bundle over x. In other words, we can view $\mathcal{H}$ as a flat real vector bundle over $\mathcal{M}_g$, equipped with a complex structure J which varies real analytically with the basepoint. In particular, we can view the Gauss-Manin connection on $H^1(S,\mathbb{R})$ as a connection on $\mathcal{H}$.

Denote by $\mathcal{H}_+ \subset \mathcal{H}$ the complement of the zero section in the Hodge bundle $\mathcal{H}$. This is a complex orbifold. The pull-back

$$\Pi^*\mathcal{H} \to \mathcal{H}_+$$

to $\mathcal{H}_+$ of the Hodge bundle on $\mathcal{M}_g$ is a holomorphic vector bundle on $\mathcal{H}_+$. The pull-back of the Gauss-Manin connection is a flat connection on $\Pi^*\mathcal{H}$. In the sequel we simply write $\mathcal{H}$ for the bundle $\Pi^*\mathcal{H}$ whenever what is meant is clear from the context.

Let $\mathcal{Q}_+ \subset \mathcal{H}_+$ be a component of a stratum. We use the notation $\mathcal{Q}_+$ to indicate that we do not normalize the area of an abelian differential. Then $\mathcal{Q}_+$ is a complex suborbifold of $\mathcal{H}_+$. Period coordinates for $\mathcal{Q}_+$ define the complex structure.

The group $GL(2,\mathbb{R})$ acts on $\mathcal{H}$, and this action preserves $\mathcal{Q}_+$. An *affine invariant manifold* $\mathcal{C}_+$ in $\mathcal{Q}_+$ is the closure in $\mathcal{Q}_+$ of an orbit of the $GL(2,\mathbb{R})$-action. Such a manifold is affine in period coordinates [EMM13]. Moreover, $\mathcal{C}_+ \subset \mathcal{Q}_+$ is a complex suborbifold of $\mathcal{Q}_+$ [EMM13]. Period coordinates determine a projection

$$p : T\mathcal{C}_+ \to \mathcal{H} \oplus \overline{\mathcal{H}} = H^1(S,\mathbb{C})$$

to absolute periods (see [W14] for a clear exposition). The image $p(T\mathcal{C}_+)$ is a flat subbundle of the flat complex vector bundle $\mathcal{H}\oplus\overline{\mathcal{H}}|_{\mathcal{C}_+} = H^1(S,\mathbb{C})|_{\mathcal{C}_+}$. The complex structure is the natural complex structure in period coordinates.

By the main result of [F13], there is a holomorphic subbundle $\mathcal{Z}$ of $\mathcal{H}|_{\mathcal{C}_+}$ such that

$$p(T\mathcal{C}_+) = \mathcal{Z} \oplus \overline{\mathcal{Z}}.$$

We call $\mathcal{Z}$ the *absolute holomorphic tangent bundle* of $\mathcal{C}_+$. As a real vector bundle, $\mathcal{Z}$ can be identified with $p(T\mathcal{C}_+) \cap H^1(S,\mathbb{R})$, and this bundle is invariant under the compatible complex structure. As a consequence, $\mathcal{Z} \subset H^1(S,\mathbb{R})|_{\mathcal{C}_+}$ is symplectic, moreover it is flat, i.e. it is invariant under the restriction of the Gauss Manin connection to $\mathcal{C}_+$.

Define the *rank* of the affine invariant manifold $\mathcal{C}_+$ as

$$\mathrm{rk}(\mathcal{C}_+) = \frac{1}{2}\dim_{\mathbb{C}} p(T\mathcal{C}_+) = \dim_{\mathbb{C}}\mathcal{Z}.$$

With this definition, components of strata are affine invariant manifolds of rank g. The *monodromy group* of $\mathcal{Z}$ for the restriction of the Gauss Manin connection to $\mathcal{C}_+$ then is a subgroup of $Sp(2\ell,\mathbb{R})$ where $\ell = \mathrm{rk}(\mathcal{C})$.

Example 4.1. If $\mathcal{C}_+$ is a Teichmüller curve then the monodromy group of $\mathcal{C}_+$ is just the Veech group of $\mathcal{C}_+$. Thus this monodromy group is a lattice in $Sp(2,\mathbb{R}) = SL(2,\mathbb{R})$, in particular it is Zariski dense in $SL(2,\mathbb{R})$.

The following result is the analog of the first part of Corollary 3.5 for affine invariant manifolds. It is a consequence of work of Wright [W15]. For its formulation, as in Section 3 we define an affine invariant manifold $\mathcal{C}_+$ of rank ℓ to be *locally Zariski dense* if for every open contractible subset U of $\mathcal{C}_+$ the subgroup of $Sp(2\ell,\mathbb{R})$ generated by the monodromy of those periodic orbits for Φ^t in $\mathcal{C}_+$ which pass through U (as introduced in Section 3) is Zariski dense in $Sp(2\ell,\mathbb{R})$. Note that this makes sense because the bundle $\mathcal{Z} \to \mathcal{C}_+$ is flat and therefore admits a family of canonical symplectic trivializations over the contractible set U. Replacing one of these trivializations by another one changes the local monodromoy group by a conjugation.

Lemma 4.2. *An affine invariant manifold is locally Zariski dense.*

Proof. Let $\mathcal{C}_+$ be an affine invariant manifold of rank $\ell \geq 1$, let $\mathcal{C} \subset \mathcal{C}_+$ be its subset of differentials of area one and let $\tilde{\mathcal{C}}$ be a component of the preimage of $\mathcal{C}$ in the Teichmüller space of marked abelian differentials.

Let (X,ω) be a translation surface (i.e. a Riemann surface X with a holomorphic one-form ω which defines a flat metric on X) containing a collection $\mathcal{Y}$ of vertical cylinders. Following [W15] we define the *cylinder shear* $u_t^{\mathcal{Y}}(X)$ to be the surface obtained from X by applying the horocycle flow to $\mathcal{Y}$ but not to the rest of X (for reasons that will be clear later we use vertical cylinders rather than horizontal cylinders as in [W15], and we use the horocycle flow which fixes the vertical direction).

By Theorem 1.10 of [W15], there is a translation surface $(X,\omega) \in \mathcal{C}$ with the following properties. Let $(\tilde{X},\tilde{\omega}) \in \tilde{\mathcal{C}}$ be a preimage of (X,ω). There is a cylinder decomposition of $\tilde{X}$ into ℓ collections $\mathcal{Y}_1,\dots,\mathcal{Y}_\ell$ of non-separating cylinders, and for each i and all t, the cylinder shear $u_t^{\mathcal{Y}_i}(\tilde{X})$ is contained in $\tilde{\mathcal{C}}$.

For each i, the collection $\mathcal{Y}_i$ consists of $r_i \geq 1$ cylinders with moduli $m_1,\dots,m_{r_i}$. By assumption, the cylinder shear $u_t^{\mathcal{Y}_i}(X)$ remains entirely in $\mathcal{C}$. By a local version of Lemma 3.1 of [W15], applied to this cylinder shear rather than to the full horocycle flow (note that the proof of this local version is identical to the proof given in Section 3 of [W15]), if these moduli are not rationally dependent then there is a proper subcollection of $\mathcal{Y}_i$ consisting of $1 \leq s < r_i$ cylinders so that the cylinder shear for this subcollection is contained in $\mathcal{C}$. But then there are at least $\ell+1$ collections of vertical cylinders with the property that the cylinder shear for each of these collections preserves $\mathcal{C}$ (see Section 3 of [W15] for details). This violates Theorem 1.10 of [W15].

As a consequence, the moduli of the cylinders in each of the collections $\mathcal{Y}_i$ are rationally dependent. Then there are numbers $k_1,\dots,k_{r_i} \geq 1$ and a number $s(i) > 0$ such that $u_{s(i)}^{\mathcal{Y}_i}(\tilde{X})$ equals the image of $\tilde{X}$ under the Dehn multi-twist T_i with multiplicities k_u about the r_i different core curves of the cylinders in $\mathcal{Y}_i$.

We now use an argument similar to the proof of Proposition 3.4 (compare also the proof of Theorem 5.1 of [W14] for an argument along these lines). Choose a contractible neighborhood U of ω in $\mathcal{C}$ which is small enough that the bundle $\mathcal{Z}|U$ is trivial as a symplectic vector bundle, with a trivialization defined by the Gauss Manin connection (with the interpretation given as preceding this proof). Let $\Omega(\Gamma_0)$ be the subsemigroup of $\mathrm{Mod}(S)$ defined as in Section 2 and 3 from a set Γ_0 of periodic orbits for Φ^t in $\mathcal{C}$ passing through a nested family of subsets of U, and let G be the subgroup of $\mathrm{Mod}(S)$ generated by $\Omega(\Gamma_0)$.

We claim that $\Psi(G)$ contains the subgroup of $Sp(2g,\mathbb{Z})$ generated by those multi-transvections which are the images of the Dehn multitwists T_i (here as before, $\Psi : \mathrm{Mod}(S) \to Sp(2g,\mathbb{Z})$ is the natural homomorphism). To this end we use the reasoning from the proof of Proposition 3.4. Namely, the Dehn multitwist T_i stabilizes the component $\tilde{\mathcal{C}}$ of the preimage of $\mathcal{C}$. Let $\tilde{U}$ be the lift of U to $\tilde{\mathcal{C}}$ containing the lift $\tilde{\omega}$, and choose a smooth extension $\zeta : \tilde{U} \times [0,1] \to \tilde{\mathcal{C}}$ of the lift of the cylinder shear of $\tilde{\omega}$ so that for each $\tilde{q} \in \tilde{U}$, the path $t \to \zeta(\tilde{q},t)$ connects $\tilde{q}$ to $T_i\tilde{q}$.

Choose a periodic point $q \in U$ for the Teichmüller flow and let $\tilde{q} \in \tilde{U}$ be the preimage of q. Let $\varphi \in \mathrm{Mod}(S)$ be the pseudo-Anosov mapping class which connects $\tilde{q}$ to $\Phi^\sigma\tilde{q}$ for some $\sigma > 0$. Then $T_i \circ \varphi \circ T_i^{-1}$ connects $T_i\tilde{q}$ to $T_i(\Phi^\sigma\tilde{q})$.

Following the proof of Proposition 5.4 of [H13], let $W^{su}_{\tilde{\mathcal{C}}}(\tilde{q})$ be the local strong unstable manifold of $\tilde{\mathcal{C}}$ containing $\tilde{q}$ in $\tilde{\mathcal{C}}$. This is a closed subset of the set of all abelian differentials in $\tilde{\mathcal{C}}$ whose horizontal measured geodesic lamination coincides with the horizontal geodesic lamination of $\tilde{q}$. The set of points in $W^{su}_{\tilde{\mathcal{C}}}(\tilde{q})$ whose forward Φ^t-orbit is contained in a fixed compact subset of $\mathcal{C}$ projects homeomorphically into the boundary of the *curve graph* of S. We may assume that this compact set contains the endpoints of both the orbit through $\tilde{q}$ as well as the orbit through $T_i\tilde{q}$. For sufficiently large k, the map φ^k acts as a contraction on this set, and the same holds true for $T_i \circ \varphi^k \circ T_i^{-1}$. The arguments in the proof of Proposition 5.4 of [H13] then show that for sufficiently large k, the periodic orbit for the pseudo-Anosov element $\varphi^k \circ T_i \circ \varphi^k$ is contained in $\mathcal{C}$. By the argument in the proof of Proposition 3.4 we now conclude that $T_i \in G$.

Now the families of cylinders $\mathcal{Y}_i$ are pairwise disjoint and therefore the subgroup of $\mathrm{Mod}(S)$ generated by the Dehn-multitwists T_i is an abelian group of rank ℓ. By Theorem 1.10 of [W15], their common fixed set intersects $\mathcal{Z} = p(T\mathcal{C}_+) \cap H^1(S,\mathbb{R})$ in a Lagrangian linear subspace A_1 of $\mathcal{Z}$.

Let again U be a small contractible neighborhood of ω with lift $\tilde{U}$ to $\tilde{\mathcal{C}}$ so that the bundle $\mathcal{Z}|U$ is trivial. For $\tilde{q} \in \tilde{U}$ the imaginary part $\mathrm{Im}(\tilde{q})$ of $\tilde{q}$ is a harmonic one-form for the complex structure underlying $\tilde{q}$ which defines a cohomology class $[\mathrm{Im}(\tilde{q})] \in \mathcal{Z} \subset H^1(S,\mathbb{R})$. Since $\mathcal{C}$ is defined by polynomial equations in period coordinates [EMM13], as $\tilde{q}$ varies in $\tilde{U}$ these cohomology classes vary through an open subset of $\mathcal{Z} \sim \mathbb{R}^{2\ell}$.

Denote by $[c_i]$ the homology class of the oriented weighted multicurve which determines the Dehn multi-twist T_i. The evaluation of the imaginary part of $\tilde{\omega}$ on each of the classes $[c_i]$ is positive. As this is an open condition, by decreasing the

size of U we may assume that $\langle [\mathrm{Im}(\tilde{q})], [c_i]\rangle \neq 0$ for all i and all $\tilde{q} \in \tilde{U}$ where $\langle , \rangle$ is the natural pairing. Now periodic orbits for Φ^t are dense in $\mathcal{C}$ and therefore we can find a periodic point $z \in U$ with preimage $\tilde{z} \in \tilde{U}$ such that $\langle [\mathrm{Im}(\tilde{z})], [c_i]\rangle \neq 0$ for all i.

Let $\varphi \in \mathrm{Mod}(S)$ be the pseudo Anosov element whose axis passes through $\tilde{z}$. There is a number $\kappa > 1$ such that $\varphi^* \mathrm{Im}(\tilde{z}) = \kappa^{-1}\mathrm{Im}(\tilde{z})$, moreover κ is the Perron Frobenius eigenvalue for the action of φ on $H_1(S, \mathbb{R})$. By invariance of the natural pairing $\langle , \rangle$ under φ, as $k \to \infty$ the homology classes $\varphi^k([c_i])$ converge up to rescaling to a class in $H_1(S, \mathbb{R})$ whose contraction with the intersection form defines $\pm[\mathrm{Im}(\tilde{z})]$, viewed as a linear functional on $H_1(S, \mathbb{R})$.

As a consequence, for suffiently large j and all i, ℓ we have $\iota([\varphi^j c_i], [c_\ell]) \neq 0$. It now follows from the arguments in the proof of Proposition 3.4 and equivariance that the group generated by $\Psi\Omega(\Gamma_0)$ contains a subgroup generated by at least $\ell+1$ multi-transvections with integral homology classes which are independent over $\mathbb{R}$. These are the images under Ψ of the Dehn multitwists T_i $(i \leq \ell)$ and images under Ψ of the Dehn multitwists $\varphi^j T_i \varphi^{-j}$. Moreover, $\iota([\varphi^j c_i], [c_u]) \neq 0$ for all i, u.

Let $A_2 \subset A_1$ be the common fixed set in $\mathcal{Z}$ of the transvections which are the images of the multitwists $T_i, \varphi^j T_u \varphi^{-j}$. Then A_2 is a linear subspace of A_1, and for large enough j its codimension in A_1 is $s \geq 1$. Let $i_1, \ldots, i_s \subset \{1, \ldots, \ell\}$ be such that the homology classes $[c_i], [\varphi^j c_{i_u}]$ $(i \leq \ell, u \leq s)$ are independent over $\mathbb{R}$ and their common fixed set in $\mathcal{Z}$ is A_2. Using again the fact that the set of imaginary parts of differentials in $\tilde{U}$ define an open subset of $p(T\mathcal{C}_+) \cap H^1(S, \mathbb{R})$, we can find some $\bar{z} \in \tilde{U}$ and some $a \in A_2$ so that $\langle [\mathrm{Im}(\bar{z})], a\rangle > 0$. As before, we may assume that $\bar{z}$ is the preimage of a periodic point. Argue now as in the previous paragraph and find a multitwist β in the group generated by $\Omega(\Gamma_0)$ so that the invariant subspace of the subgroup generated by $\Psi(\beta)$ and A_2 has codimension at least one in A_2.

Repeat this construction. In finitely many steps we find integral homology classes $a_1, \ldots, a_\ell, a_{\ell+1}, \ldots, a_{2\ell} \in H_1(S, \mathbb{R})$ (where for $i \leq \ell$ the class a_i is the class of the weighted multicurve which determines T_i) with the following properties.

(1) Let $V \subset H_1(S, \mathbb{R})$ be the real vector space spanned by the classes a_i. Then the dimension of V equals 2ℓ and the restriction of V to $\mathcal{Z}$ is non-degenerate (where we view V as a subspace of $H^1(S, \mathbb{R})^*$). In particular, V is a symplectic subspace of $H_1(S, \mathbb{R})$.
(2) $\iota(a_{\ell+1}, a_i) \neq 0$ for all $i \leq \ell$, and for all $j \geq \ell + 2$ there is some $i \leq \ell$ such that $\iota(a_j, a_i) \neq 0$.
(3) For each i the multi-transvection $b \to b + \iota(b, a_i)a_i$ is contained in the group generated by $\Psi(\Omega(\Gamma_0))$.

By the choice of the homology classes a_i, the $(2\ell, 2\ell)$-matrix $(\iota(a_i, a_j))$ whose (i, j)-entry is the intersection $\iota(a_i, a_j)$ is integral and of maximal rank. Choose a prime $p \geq 5$ so that each of the entries of $(\iota(a_i, a_j))$ is prime to p. All but finitely many primes will do. Then the reduction mod p of the matrix $(\iota(a_i, a_j))$ is of maximal rank as well. In particular, if F_p denotes the field with p elements then

the reductions mod p of the homology classes a_i span a 2ℓ-dimensional symplectic subspace V_p of $H_1(S, F_p)$.

Let $G < \mathrm{Sp}(V)$ be the subgroup of the sympelctic group of V which is generated by the multi-transvections with the elements a_i. Its reduction G_p mod p preserves the symplectic subspace V_p of $H_1(S, F_p)$ and acts on V_p as a group of symplectic transformations. Lemma 3.2 shows that $G_p = Sp(2\ell, F_p)$. Note that property (2) above guarantees that all conditions in Lemma 3.2 are fulfilled. Then G is a Zariski dense subgroup of the group of symplectic automorphisms of V [Lu99]. By duality and the discussion in the proof of Proposition 3.4, this just means that $\mathcal{C}_+$ is locally Zariski dense. □

Recall that the pull-back $\Pi^*\mathcal{H} \to \mathcal{H}_+$ to $\mathcal{H}_+$ of the Hodge bundle on $\mathcal{M}_g$ is a holomorphic vector bundle on $\mathcal{H}_+$. The Hermitean structure on $\mathcal{H}$ defines a Hermitean structure on $\Pi^*\mathcal{H}$. The bundle $\Pi^*\mathcal{H}$ naturally splits as a direct sum

$$\Pi^*\mathcal{H} = \mathcal{T} \oplus \mathcal{L}$$

of complex vector bundles. Here the fibre of $\mathcal{T}$ over a point q is just the $\mathbb{C}$-span of q, and the fibre of $\mathcal{L}$ is the orthogonal complement of $\mathcal{T}$ for the natural Hermitean metric. The complex line bundle $\mathcal{T}$ is holomorphic. Via identification of $\mathcal{L}$ with the quotient bundle $\Pi^*\mathcal{H}/\mathcal{T}$, we may assume that $\mathcal{L}$ is holomorphic. Its complex dimension equals $g - 1 \geq 2$.

For each $q \in \mathcal{H}_+$, the $GL(2, \mathbb{R})$-orbit through q locally preserves the two-dimensional symplectic subspace of $H^1(S, \mathbb{R})$ spanned by the real and imaginary part of q. This implies that the restriction of the Gauss Manin connection to an orbit of the action of $GL(2, \mathbb{R})$ induces a flat connection ∇^{GM} on the symplectic complement of the $\mathbb{C}$-span of q in $\mathcal{H}$, where this $\mathbb{C}$-span is taken with respect to the complex structure J_q.

With respect to the above identification of $\mathcal{H}$ with $H^1(S, \mathbb{R})$, this just means that the Gauss-Manin connection restricts to a flat leafwise connection ∇^{GM} on the bundle $\mathcal{L}$ over the foliation of $\mathcal{H}_+$ into the orbits of the $GL(2, \mathbb{R})$-action. Here a leafwise connection is a connection where covariant derivative is only defined for vectors tangent to the foliation. As the Gauss Manin connection is real analytic in period coordinates, the $GL(2, \mathbb{R})$-action is real analytic and the splitting $\Pi^*\mathcal{H} = \mathcal{T} \oplus \mathcal{L}$ is real analytic, the leafwise connection is real analytic. It preserves the symplectic structure of $\mathcal{L}$, but there is no information on the complex structure. In particular, parallel transport for ∇^{GM} defines an extension Θ^t of the Teichmüller flow Φ^t on $\mathcal{H}_+$ to a flow on $\mathcal{L}$ which preserves the symplectic structure of $\mathcal{L}$. Moreover, for each $k \leq g - 2$ the leafwise connection ∇^{GM} extends to a flat leafwise connection on the bundle $\wedge^{2k}_{\mathbb{R}}\mathcal{L}$ (whose fibre at q is the $2k$-th exterior power of the fibre of $\mathcal{L}$ at q, viewed as a real vector space).

The Hermitean metric on the holomorphic vector bundle $\Pi^*\mathcal{H}$ determines a unique Chern connection ∇ on $\Pi^*\mathcal{H}$. Viewing again $\Pi^*\mathcal{H}$ as the real vector bundle $H^1(S, \mathbb{R})$ with the complex structure J, the complex structure J is parallel for ∇. Since the $GL(2, \mathbb{R})$-orbits on $\mathcal{H}_+$ are holomorphic suborbifolds of $\mathcal{H}_+$, by naturality the restriction of the Chern connection to $GL(2, \mathbb{R})$-orbits preserves the decomposition $\Pi^*\mathcal{H} = \mathcal{T} \oplus \mathcal{L}$.

As a consequence, parallel transport for ∇ defines an extension of the Teichmüller flow Φ^t on $\mathcal{H}_+$ to a flow Ψ^t on $\mathcal{L}$. This extension commutes with the complex structure J, viewed as a bundle automorphism whose square equals $-\mathrm{Id}$. Similarly, for every $k \leq g-2$ the complex structure J extends to an automorphism of the real tensor bundle $\wedge^{2k}_{\mathbb{R}}\mathcal{L}$, and the restriction of the connection ∇ to the orbits of the $GL(2,\mathbb{R})$-action extends to a connection on $\wedge^{2k}_{\mathbb{R}}\mathcal{L}$ which commutes with this automorphism.

The Hermitean metric which determines the Chern connection is defined by the polarization and the complex structure. These data are real analytic in period coordinates (recall that the Torelli map is holomorphic) and consequently the connection matrix for the Chern connection in period coordinates is real analytic (see [GH78]).

To summarize, for every $k \leq g-2$, both the Chern connection and the Gauss Manin connection restrict to leafwise connections of the restriction of the bundles $\wedge^{2k}_{\mathbb{R}}\mathcal{L} \to \mathcal{H}_+$ to the orbits of the $GL(2,\mathbb{R})$-action, and these leafwise connections depend in a real analytic fashion on period coordinates.

The foliation of $\mathcal{H}_+$ into the orbits of the $GL(2,\mathbb{R})$-action is real analytic and hence has a tangent bundle $\Sigma \subset T\mathcal{H}_+$ (which can naturally be identified with the bundle $\mathcal{T}$). Then Σ is a real analytic subbundle of $T\mathcal{H}_+$ which is naturally trivialized by the generators X, Y, Z of the Teichmüller flow and the positive and negative horocycle flow as well as be the generator of the center of $GL(2,\mathbb{R})$. By the above discussion, $\nabla - \nabla^{GM}$ defines a real analytic tensor field

$$\Xi \in \Omega(\Sigma^* \otimes \mathcal{L}^* \otimes \mathcal{L})$$

where we denote by $\Omega(\Sigma^* \otimes \mathcal{L}^* \otimes \mathcal{L})$ the vector space of real analytic sections of the real analytic vector bundle $\Sigma^* \otimes \mathcal{L}^* \otimes \mathcal{L}$. For every $k \leq g-2$ the tensor field Ξ^k defined as the action of $\nabla - \nabla^{GM}$ on $\wedge^{2k}_{\mathbb{R}}\mathcal{L}$ is real analytic as well. If $\mathcal{C}_+ \subset \mathcal{H}_+$ is an affine invariant manifold of rank $2 \leq \ell \leq g-1$, with absolute holomorphic tangent bundle $\mathcal{Z}$, then the restriction of the tensor field $\Xi^{\ell-1}$ to $\mathcal{C}_+$ preserves up to scale the J-invariant section of $\wedge^{2\ell-2}_{\mathbb{R}}\mathcal{L}$ which is defined by $p(T\mathcal{C}_+) \cap H^1(S,\mathbb{R})$.

The following proposition is the key step towards the proof of Theorem 3 from the introduction.

Proposition 4.3. *Let $\mathcal{C}_+ \subset \mathcal{Q}_+$ be an affine invariant manifold of rank $\ell \geq 3$. There are finitely many proper affine invariant submanifolds of $\mathcal{C}_+$ which contain every affine invariant submanifold of $\mathcal{C}_+$ of rank $2 \leq k \leq \ell - 1$.*

Proof. Let $\mathcal{Q}_+ \subset \mathcal{H}_+$ be a component of a stratum and let $\mathcal{C}_+ \subset \mathcal{Q}_+$ be an affine invariant manifold, with absolute holomorphic tangent bundle $\mathcal{Z} \to \mathcal{C}_+$. As before, there is a splitting

$$\mathcal{Z} = \mathcal{T} \oplus (\mathcal{L} \cap \mathcal{Z}).$$

The bundle

$$\mathcal{W} = \mathcal{L} \cap \mathcal{Z} \to \mathcal{C}_+$$

is holomorphic. It also can be viewed as a real analytic real vector bundle with a real analytic complex structure J (which is just a real analytic section of the real analytic endomorphism bundle of $\mathcal{W}$ with $J^2 = -\mathrm{Id}$).

For $1 \leq k \leq \ell - 2$ denote by $\mathrm{Gr}(2k) \to \mathcal{C}_+$ the fibre bundle whose fibre over ω is the Grassmannian of oriented $2k$-dimensional real subspaces of $\mathcal{W}_\omega$. This is a real analytic fibre bundle with compact fibre. It contains the bundle $\mathcal{P}(k) \to \mathcal{C}_+$ whose fibre over ω is the Grassmannian of *complex* k-dimensional subspaces of $\mathcal{W}_\omega$ (for the complex structure J) as a real analytic subbundle.

The Hermitean metric on $\mathcal{W}$ naturally extends to a real analytic Riemannian metric on $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$. The bundle $\mathrm{Gr}(2k)$ can be identified with the set of indecomposable points in the sphere subbundle of $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$ for this extension. Namely, an oriented $2k$-dimensional real subspace of $\mathcal{W}_\omega$ defines uniquely a point in $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$ of norm one which is just the exterior product of an orthonormal basis of the subspace with respect to the metric on $\mathcal{W}$. The points in $\mathcal{P}(k)$ correspond precisely to those indecomposable points which are invariant under the extension of J to an automorphism of $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$.

Let B^k (or C^k, D^k) be the contraction of the tensor field Ξ^k with the vector fields X, Y, Z. Since these vector fields are real analytic, B^k (or C^k, D^k) is a real analytic section of the endomorphism bundle $\mathrm{End}(\wedge^{2k}_{\mathbb{R}}(\mathcal{W}))$.

Let again $\mathcal{C} \subset \mathcal{C}_+$ be the real analytic hyperplane of abelian differentials in $\mathcal{C}$ of area one. For $1 \leq k \leq \ell - 2$ and $\omega \in \mathcal{C}$ let

$$\rho^k(\omega, 0) \subset \mathcal{P}(k)_\omega$$

be the set of all complex linear subspaces L with $B^k L \subset L, C^k L \subset L, D^k L \subset L$. By this we mean that $L \in \rho^k(\omega, 0)$ if the point in $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$ defined by L is mapped by B^k, C^k, D^k to a multiple of itself. (Strictly speaking, we do not need the endomorphisms C^k, D^k for our argument, but we introduced them to make the argument more transparent).

Define a *real analytic subset* of $\mathcal{P}(k)$ to be the intersection of the zero sets of a finite or countable number of real analytic functions on $\mathcal{P}(k)$. Recall that this is well defined since $\mathcal{P}(k)$ has a natural real analytic structure. We allow such functions to be constant zero, i.e. we do not exclude that such a set coincides with $\mathcal{P}(k)$. The real analytic set is *proper* if it does not coincide with $\mathcal{P}(k)$. Then there is at least one defining function which is not identically zero and hence the set is closed and nowhere dense.

We claim that $\cup_\omega \rho^k(\omega, 0)$ is a real analytic subset of $\mathcal{P}(k)$. To this end observe that $x \in \rho^k(\omega, 0)$ if the unit vector $Z \in \wedge^{2k}_{\mathbb{R}}\mathcal{W}$ corresponding to x satisfies $Z \wedge B^k Z = 0 = Z \wedge C^k Z = Z \wedge D^k Z$ (here the wedge product is taken in $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$). As B^k, C^k, D^k are real analytic tensor field, $Z \to Z \wedge B^k Z$ is a real analytic bundle map and similarly for C^k, D^k. Hence the preimage of the zero section of the vector bundle $(\wedge^{2k}_{\mathbb{R}}\mathcal{W}) \wedge (\wedge^{2k}_{\mathbb{R}}\mathcal{W})$ is a real analytic subset of the real analytic set of points in the sphere bundle of $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$ which correspond to points in $\mathcal{P}(k)$.

For $t > 0$ define $\rho^k(\omega, t) \subset \mathrm{Gr}(2k)_\omega$ to be the preimage of $\rho^k(\Phi^t\omega, 0)$ under parallel transport for the Gauss Manin connection along the flow line $s \to \Phi^s\omega$ ($s \in [0, t]$). By the discussion in the previous paragraph and the fact that parallel

transport is real analytic, $\cup_\omega \rho^k(\omega,t)$ is a real analytic subset of $\mathrm{Gr}(2k)_\omega$ and hence the same holds true for

$$\xi^k = \cap_{t\in\mathbb{R}} \cup_\omega \rho^k(\omega,t)$$

(take the intersections over all rationals in $\mathbb{R}$). By construction, the set ξ^k is invariant under the extension Θ_k^t of the Teichmüller flow by parallel transport of fibres for the Gauss Manin connection. If $\mathcal{U} \subset \mathcal{C}$ is a proper affine invariant manifold of rank $k+1 < \ell$ then it follows from the discussion preceding this proof (see [F13]) that for every $\omega \in \mathcal{U}$ the projected tangent space $p(T\mathcal{U})$ defines a point in $\xi^k(\omega) \subset \mathcal{P}(k)_\omega$.

Let $\pi : \mathcal{P}(k) \to \mathcal{C}$ be the natural projection and let

$$\mathcal{A} = \pi(\xi^k).$$

Since the projection π is closed, $\mathcal{A}$ is a closed Φ^t-invariant subset of $\mathcal{C}$ which contains all affine invariant submanifolds of $\mathcal{C}$ of rank $k+1$. We claim that $\mathcal{A}$ is nowhere dense in $\mathcal{C}$.

Assume to the contrary that $\mathcal{A}$ contains an open subset U of $\mathcal{C}$. Then there is for every $q \in U$ a line in $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$ spanned by an indecomposable vector Z which is an eigenvector for the extension of J and which also is an eigenvector for B^k, C^k, D^k for some real eigenvalue. Moreover, the same holds true for the parallel transport for the Gauss Manin connection of this eigenline along the flow line for Φ^t through q. Since the operators J, B^k, C^k, D^k are linear, we conclude that for each $q \in U$ there is a non-trivial linear subspace $E(q) = \sum_i E_i(q) \subset \wedge^{2k}_{\mathbb{R}}\mathcal{W}_q$ with the following properties.

(1) $E(q)$ is the linear span of $\xi^k(q)$, in particular $E(q)$ is contained in the eigenspace of J for the eigenvalue one, and it is spanned by indecomposable vectors.
(2) Each vector space $E_i(q)$ is spanned by indecomposable vectors.
(3) For each i, the parallel transport of a vector in the subspace $E_i(q)$ along a flow line of the Teichmüller flow with respect to the Chern connection coincides up to scale with its parallel transport for the Gauss Manin connection.

As ξ^k is real analytic and invariant under the extension Θ_k^t of the Teichmüller flow via parallel transport for the Chern connection, there is an open Φ^t-invariant subset V of U so that E and its decomposition depend continuously on $q \in V$. In particular, the dimensions of the subspaces $E_i(q)$ are constant on V.

Assume first that for $q \in V$ the rank of $E(q)$ is strictly smaller than the rank of the eigenspace of J in $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$ for the eigenvalue one. This implies that there is a proper analytic subbundle of

$$\mathcal{P}(k) \to \cup_{t\in\mathbb{R}} \Phi^t V$$

whose fibre over a point $q \in V$ is an intersection of hyperplanes in $\mathcal{P}(k)_q$ and which is invariant under parallel transport for the Gauss-Manin connection. But then the local monodromy group (as defined in the beginning of this section) of a small contractible open subset of V is not Zariski dense in $Sp(2\ell,\mathbb{R})$ contradicting Lemma

4.2. The same argument also rules out the possibility that the decomposition $E = \sum_i E_i(q)$ is non-trivial.

Finally assume that E coincides with the eigenspace for the action of J on $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$. This then implies that the tensor field Ξ^k maps the automorphism J of $\wedge^{2k}_{\mathbb{R}}\mathcal{W}$ to a real multiple. But then the local monodromy of a small contractible subset of V preserves k-dimensional complex subspaces of $\mathcal{W}$ which again violates Lemma 4.2.

As a consequence, the set $\mathcal{A}$ is closed and nowhere dense, and it contains all affine invariant submanifolds of $\mathcal{C}$ of rank $k+1$. Thus all affine invariant submanifolds of $\mathcal{C}$ are contained in the complement of the saturation of $\mathcal{C}-\mathcal{A}$ for the $SL(2,\mathbb{R})$-action. This complement is a closed nowhere dense $SL(2,\mathbb{R})$-invariant subset of $\mathcal{C}$. By Theorem 2.2 of [EMM13], we conclude that indeed, there are finitely many proper affine invariant submanifolds of $\mathcal{C}$ which contain all affine invariant submanifolds of rank $k+1$. This completes the proof of the proposition. □

As an immediate corollary, we obtain Theorem 3 from the introduction.

Corollary 4.4. (1) *Let $\mathcal{Q}$ be a component of a stratum; then for every $2 \leq \ell \leq g-1$ there are finitely many affine invariant submanifolds of $\mathcal{Q}$ of rank ℓ which contain every affine invariant submanifold of rank ℓ.*

(2) *The smallest stratum of differentials with a single zero contains only finitely many affine invariant submanifolds of rank at least two.*

Proof. To show the first part of the corollary, let $2 \leq \ell \leq g-1$. By Proposition 4.3, applied to $\mathcal{Q}$, there are finitely many proper affine invariant submanifolds $\mathcal{C}_1, \dots, \mathcal{C}_k$ of $\mathcal{Q}$ which contain every affine invariant manifold of rank ℓ. The dimension of $\mathcal{C}_i$ is strictly smaller than the dimension of $\mathcal{Q}$.

Among the submanifolds $\mathcal{C}_i$, assume that for all $i \leq u$ the rank of $\mathcal{C}_i$ is bigger than ℓ and that the rank of $\mathcal{C}_i$ for $i > u$ is at most ℓ. Apply Proposition 4.3 to each of the affine invariant manifolds $\mathcal{C}_i$ ($i \leq u$). In finitely many steps we deduce the statement of the first part of the corollary.

Now let $\mathcal{H}(2g-2)$ be a stratum of differentials with a single zero. Period coordinates for $\mathcal{H}(2g-2)$ are given by absolute periods, and the dimension of an affine invariant manifold $\mathcal{C} \subset \mathcal{H}(2g-2)$ of rank ℓ equals 2ℓ. Thus $\mathcal{C}$ does not contain any proper affine invariant submanifold of rank ℓ.

By Proposition 4.3, there are finitely many proper affine invariant manifolds $\mathcal{C}_1, \dots, \mathcal{C}_s$ which contain every affine invariant submanifold of $\mathcal{H}(2g-2)$ of rank $\ell \in [2, g-1]$. Then each affine invariant manifold of rank $g-1$ is contained in this collection, in particular there are only finitely many such manifolds.

To show finiteness of affine invariant manifolds of any rank $2 \leq \ell \leq g-1$, apply Proposition 4.3 to each of the manifolds $\mathcal{C}_i$ and proceed by inverse induction on the rank. □

5. Algebraically primitive Teichmüller curves

For a translation surface (X, ω) whose Veech group contains a pseudo-Anosov element, the *holonomy field* of (X, ω) coincides with the trace field of the pseudo-Anosov element (see e.g. the appendix of [KS00]- we do not need any additional information on this field). By Lemma 2.10 of [LNW15], if $\mathcal{C}$ is a rank one affine invariant manifold then for all $(X, \omega) \in \mathcal{C}$, the holonomy field of (X, ω) equals the *field of definition* of $\mathcal{C}$ [W14].

We have now a closer look at rank one affine invariant manifolds $\mathcal{C}$ whose field of definition $\mathfrak{k}$ is of degree g over $\mathbb{Q}$. Then $\mathfrak{k}$ is a totally real [F13] number field of degree g, with ring of integers $\mathcal{O}_{\mathfrak{k}}$. Via the g field embeddings $\mathfrak{k} \to \mathbb{R}$, the group $SL(2, \mathcal{O}_{\mathfrak{k}})$ embeds into $G = SL(2, \mathbb{R}) \times \cdots \times SL(2, \mathbb{R}) < Sp(2g, \mathbb{R})$ and in fact, $SL(2, \mathcal{O}_{\mathfrak{k}})$ is a lattice in G. The trace field of every periodic orbit γ in $\mathcal{C}$ equals $\mathfrak{k}$ and hence the corresponding pseudo-Anosov element $A(\gamma)$ maps with the homomorphism $\Psi : \mathrm{Mod}(S) \to Sp(2g, \mathbb{R})$ to an element in $SL(2, \mathcal{O}_{\mathfrak{k}})$.

The following observation is immediate from Lemma 4.2 and [G12]. For its formulation, define the *extended local monodromy group* of an open contractible subset U of $\mathcal{C}$ to be the subgroup of $SL(2, \mathcal{O}_{\mathfrak{k}})$ which is generated by the monodromy of those (parametrized) periodic orbits for Φ^t in $\mathcal{C}$ which pass through U.

Lemma 5.1. *For a rank one affine invariant manifold $\mathcal{C}$ whose field of definition is of degree g over $\mathbb{Q}$, any extended local monodromoy group is Zariski dense in $SL(2, \mathbb{R}) \times \cdots \times SL(2, \mathbb{R})$.*

Proof. By Lemma 4.2, the projection of the extended local monodromy group of an open set $U \subset \mathcal{C}$ to the first factor $SL(2, \mathbb{R})$ of $G = SL(2, \mathbb{R}) \times \cdots \times SL(2, \mathbb{R})$ is Zariski dense in $SL(2, \mathbb{R})$ and hence it is non-elementary. Moreover, by definition and [KS00, LNW15], the invariant trace field of the extended local monodromy group equals $\mathfrak{k}$ (compare [G12] for the definition of the invariant trace field). Thus by Corollary 2.2 of [G12], the extended local monodromy group of U is Zariski dense in G. □

In the statement of the next corollary, the affine invariant manifold $\mathcal{B}_+$ may be a component of a stratum.

Corollary 5.2. *Let $\mathcal{C}_+$ be a rank one affine invariant manifold whose field of definition is of degree g over $\mathbb{Q}$. Assume that $\mathcal{C}_+$ is properly contained in an affine invariant manifold $\mathcal{B}_+$ of rank at least three. Let $\mathcal{Z} \to \mathcal{B}_+$ be the absolute holomorphic tangent bundle of $\mathcal{B}_+$; then $\mathcal{Z}|\mathcal{C}_+$ splits as a sum of holomorphic line bundles which are invariant under both the Chern connection and the Gauss Manin connection.*

Proof. In the case that the rank of $\mathcal{B}_+$ equals g (and hence $\mathcal{Z} = \mathcal{H}$), the statement is immediate from Theorem 1.5 of [W14]. Thus assume that the rank of $\mathcal{B}_+$ is at most $g - 1$.

Since $\mathcal{C}_+ \subset \mathcal{B}_+$, the restriction of $\mathcal{H}$ to $\mathcal{C}_+$ has two splittings which are invariant under the extended local monodromy of $\mathcal{C}_+$. The first splitting is the splitting into g

line bundles obtained from the different field embddings of the field of definition of $\mathcal{C}_+$ into $\mathbb{R}$ (see Theorem 1.5 of [W14]). The second splitting is the splitting into the absolute holomorphic tangent bundle $\mathcal{Z}$ of $\mathcal{B}_+$ (which is a holomorphic subbundle of $\mathcal{H}|_{\mathcal{B}_+}$ whose complex rank equals the rank of $\mathcal{B}_+$) and its symplectic complement. Since by Lemma 5.1 the extended local monodromy group of $\mathcal{C}_+$ is Zariski dense in $SL(2,\mathbb{R}) \times \cdots \times SL(2,\mathbb{R})$, the bundle $\mathcal{Z}|\mathcal{C}_+$ is a sum of invariant line bundles. □

Corollary 5.3. *For a component $\mathcal{Q}$ of a stratum in genus $g \geq 3$, all affine invariant submanifolds of rank one whose field of definition is of degree g over $\mathbb{Q}$ are contained in a finite collection of affine invariant submanifolds of rank at most two.*

Proof. Let $\mathfrak{C}$ be the collection of all rank one affine invariant submanifolds of $\mathcal{Q}$ whose field of definition is a number field of degree g over $\mathbb{Q}$. For each $\mathcal{C}_+ \in \mathfrak{C}$, the restriction of the bundle $\mathcal{L}$ to $\mathcal{C}_+$ splits as a sum of flat holomorphic line bundles which are invariant under the Gauss-Manin connection. Thus by Proposition 4.3 and its proof, all but finitely many elements of $\mathfrak{C}$ are submanifolds of a finite set of affine invariant submanifolds of $\mathcal{Q}$ of rank at most $g-1$.

Now if $\mathcal{B}_+$ is an affine invariant manifold of rank contained in $[3, g-1]$ which contains some $\mathcal{C}_+ \in \mathfrak{C}$ then by Corollary 5.2 the above reasoning can be applied to $\mathcal{B}_+$. In finitely many steps we find finitely many proper affine invariant manifolds $\mathcal{C}_1, \dots, \mathcal{C}_k \subset \mathcal{Q}_+$ of rank at most two which contain every $\mathcal{C}_+ \in \mathfrak{C}$. This is the statement of the corollary. □

Now we are ready to complete the proof of Theorem 2.

Corollary 5.4. *For $g \geq 3$ the $SL(2,\mathbb{R})$-orbit closure of a typical periodic orbit in any component of a stratum is the entire stratum.*

Proof. Let $\mathcal{Q}$ be a component of a stratum and let $U \subset \mathcal{Q}$ be a non-empty open set. Then a typical periodic orbit for Φ^t passes through U [H13]. Thus by Proposition 4.3, the $SL(2,\mathbb{R})$-orbit closure of a typical periodic orbit either is an affine invariant manifold of rank g, or it is an affine invariant manifold of rank one.

By Theorem 1.3 of [MW15], an proper affine invariant submanifold $\mathcal{C}$ of $\mathcal{Q}$ of rank g equals the hyperelliptic locus of $\mathcal{Q}$ and $\mathcal{Q}$ is not a hyperelliptic component. Then a typical periodic orbit omits this locus. Thus we are left with ruling out that the rank of the orbit closure equals one.

By the second part of Theorem 1, the trace field of a typical perodic orbit γ is totally real and of degree g over $\mathbb{Q}$. If the rank of the $SL(2,\mathbb{R})$-orbit closure $\mathcal{C}$ of γ equals one then this trace field is the field of definition of $\mathcal{C}$ [LNW15]. Thus the corollary follows from Corollary 5.3. □

We complete this work with the proof of Theorem 4. We begin with

Proposition 5.5. *Let $g \geq 3$ and let $\mathcal{B}_+ \subset \mathcal{Q}_+$ be a rank two affine invariant manifold. Then the union of all algebraicially primitive Teichmüller curves which are contained in $\mathcal{B}_+$ does not have an accumulation point in $\mathcal{B}_+$.*

Proof. Let $\mathcal{B}_+ \subset \mathcal{Q}_+$ be a rank two affine invariant manifold. We argue by contradiction, and we assume that there is an accumulation point $\omega \in \mathcal{B}_+$ for algebraically primitive Teichmüller curves $\mathcal{C}_+ \subset \mathcal{B}_+$.

Let $\mathcal{Z} \to \mathcal{B}_+$ be the absolute holomorphic tangent bundle of $\mathcal{B}_+$. Let U be a small contractible neighborhood of ω so that there is a trivialization of the Hodge bundle over U defined by the Gauss Manin connection. The extended local monodromy group of U preserves $\mathcal{Z}$. Let $\mathcal{C}_i \subset \mathcal{B}_+$ be a sequence of algebraically primitive Teichmüller curves which accumulate at ω. Let $\mathfrak{k}_i$ be the trace field of $\mathcal{C}_i$. We claim that for all but finitely many i we have $\mathfrak{k}_i = \mathfrak{k}$ for some fixed field $\mathfrak{k}$.

Namely, assume otherwise. Then there are affine invariant manifolds $\mathcal{C}_i, \mathcal{C}_j$ which intersect U and whose fields of definition are distinct. Let G_i, G_j be the subgroups of $Sp(2g, \mathbb{R})$ isomorphic to $SL(2, \mathbb{R}) \times \cdots \times SL(2, \mathbb{R})$ which contain the groups $SL(2, \mathcal{O}_{\mathfrak{k}_i}), SL(2, \mathcal{O}_{\mathfrak{k}_j})$ as lattices. By Lemma 5.1, the extended local monodromy groups of $\mathcal{C}_i \cap U$ and $\mathcal{C}_j \cap U$ are Zariski dense in G_i, G_j. Therefore the Zariski closure in $Sp(2g, \mathbb{R})$ of the extended local monodromy group of $U \subset \mathcal{B}_+$ contains $G_i \cup G_j$. But $\mathfrak{k}_i \neq \mathfrak{k}_j$ and therefore a subgroup of $Sp(2g, \mathbb{R})$ which contains $G_i \cup G_j$ can not preserve the subspace $\mathcal{Z}$. This shows that indeed, $\mathfrak{k}_i = \mathfrak{k}$ for all but finitely many i.

Let $\mathbf{H}^2$ be the hyperbolic plane. The quotient of $\mathbf{H}^2 \times \cdots \times \mathbf{H}^2$ under the action of the group $SL(2, \mathcal{O}_{\mathfrak{k}})$ is the Hilbert modular variety $H(\mathfrak{k}) \subset \mathcal{A}_g$ of abelian varieties with real multiplication with $\mathfrak{k}$. The underlying Riemann surface of a point in one of the Teichmüller curves $\mathcal{C}_i$ is mapped by the Torelli map into $H(\mathfrak{k})$ [Mo06]. Then $\omega \in \mathcal{B}_+$ is an accumulation point in $\mathcal{B}_+$ of the set of all points which are mapped by the composition of the foot-point projection $\mathcal{B}_+ \to \mathcal{M}_g$ with the Torelli map into $H(\mathfrak{k})$. But $H(\mathfrak{k})$ is a complex submanifold of $\mathcal{A}_g$ and this composition map is holomorphic and therefore the image of $\mathcal{B}_+$ is contained in $H(\mathfrak{k})$.

As a consequence, each point in $\mathcal{B}_+$ is an abelian differential whose Jacobian has real multiplication with $\mathfrak{k}$. Moreover, the set of differentials in $\mathcal{B}_+$ which are mapped to eigenforms for this real multiplication has an acccumulation point. This implies as before that each point in $\mathcal{B}_+$ corresponds to such an eigenform and hence $\mathcal{B}_+$ is a rank one affine invariant manifold, contrary to our assumption. The proposition follows □

Proof of Theorem 4:

Let $\mathcal{Q}$ be a component of a stratum in genus $g \geq 3$. By Corollary 5.3, there are finitely many affine invariant submanifolds $\mathcal{B}_1, \ldots, \mathcal{B}_k$ of rank two which contain all but finitely many algebraically primitive Teichmüller curves.

Let $\mathcal{B}_i$ be such an affine invariant manifold of rank two. Assume that its dimension equals r for some $r \geq 4$. By Proposition 5.5, the union of all algebraically primitive Teichmüller curves which are contained in $\mathcal{B}_i$ does not have an accumulation point in $\mathcal{B}_i$. Hence this union is contained in a finite union of proper affine invariant submanifolds of $\mathcal{B}_i$ [EMM13]. The dimension of each of these invariant submanifolds is at most $r - 1$.

If there are submanifolds of rank two in this collection then we can repeat this argument to each of these finitely many submanifolds. By inverse induction on the dimension, this yields that all but finitely many algebraically primitive Teichmüller curves are contained in one of finitely many affine invariant manifolds of rank one. The field of definition of such a manifold coincides with the field of definition of the Teichmüller curve, in particular it is of degree g [LNW15].

By the main result of [LNW15], a rank one affine invariant manifold with field of definition of degree g over $\mathbb{Q}$ only contains finitely many Teichmüller curves. Thus the number of algebraically primitive Teichmüller curves in $\mathcal{Q}$ is finite as promised.

References

[Ap15] P. Apisa, *$GL_2\mathbb{R}$ orbit closures in hyperelliptic components of strata*, arXiv:1508.05438.

[ABEM12] J. Athreya, A. Bufetov, A. Eskin, M. Mirzakhani, *Lattice point asymptotic and volume growth on Teichmüller space*, Duke Math. J. 161 (2012), 1055–1111.

[Au14] D. Aulicino, *Affine manifolds and zero Lyapunov exponents in genus 3*, arXiv:1409.5180, to appear in Geom. Func. Anal.

[AEM12] A. Avila, A. Eskin, and M. Möller, *Symplectic and isometric $SL(2,\mathbb{R})$-invariant subbundles of the Hodge bundle*, arXiv:1209.2854, to appear in J. reine and angew. Math.

[AV07] A. Avila, M. Viana, *Simplicity of Lyapunov spectra: Proof of the Kontsevich-Zorich conjeccture*, Acta Math. 198 (2007), 1–56.

[BHM14] M. Bainbridge, P. Habegger, M. Möller, *Teichmüller curves in genus three and just likely intersections in $G_m^n \times G_a^n$*, arXiv:1410.6835.

[EM11] A. Eskin, M. Mirzakhani, *Counting closed geodesics in moduli space*, J. Mod. Dynamics 5 (2011), 71–105.

[EMR12] A. Eskin, M. Mirzakhani, K. Rafi, *Counting closed geodesics in strata*, arXiv:1206.5574.

[EMM13] A. Eskin, M. Mirzakhani, A. Mohammadi, *Isolation theorems for $SL(2,\mathbb{R})$-invariant submanifolds in moduli space*, arXiv:1305.3015, to appear in Ann. Math.

[FM12] B. Farb, D. Margalit, *A primer on mapping class groups*, Princeton Univ. Press, Princeton 2012.

[F13] S. Filip, *Semisimplicity and rigidity of the Kontsevich-Zorich coccyle*, arXiv:1307.7314.

[G12] S. Geninska, *Examples of infinite covolume subgroups of $PSL(2,\mathbb{R})^r$ with big limit sets*, Math. Zeit. 272 (2012), 389–404.

[GH78] P. Griffith, J. Harris, *Principles of algebraic geometry*, John Wiley & Sons, 1978.

[Hl08] C. Hall, *Big symplectic or orthogonal monodromy modulo ℓ*, Duke Math. J. 141 (2008), 179–203.

[H10] U. Hamenstädt, *Dynamics of the Teichmüller flow on compact invariant sets*, J. Mod. Dynamics 4 (2010), 393–418.

[H13] U. Hamenstädt, *Bowen's construction for the Teichmüller flow*, J. Mod. Dynamics 7 (2013), 489–526.

[H11] U. Hamenstädt, *Symbolic dynamics for the Teichmüller flow*, arXiv:1112.6107.

[H15] U. Hamenstädt, *Dynamical properties of the absolute period foliation*, preprint October 2015.

[KS00] R. Kenyon, J. Smillie, *Billiards on rational-angled triangles*, Comm. Math. Helv. 75 (2000), 65–108.

[KZ03] M. Kontsevich, A. Zorich, *Connected components of the moduli space of Abelian differentials with prescribed singularities*, Invent. Math. 153 (2003), 631–678.

[LNW15] E. Lanneau, D.-M. Nguyen, A. Wright, *Finiteness of Teichmüller curves in non-arithmetic rank 1 orbit closures*, arXiv:1504.03742.

[Lu99] A. Lubotzky, *One for almost all: generation of SL(n,p) by subsets of SL(n,Z)*, in "Algebra, K-theory, groups and education", T. Y. Lam and A. R. Magid, Editors, Contemp. Math. 243 (1999),

[MW15] C. Matheus, A. Wright, *Hodge-Teichmüller planes and finiteness results for Teichmüller curves*, Duke Math. J. 164 (2015), 1041–1077.
[M82] H. Masur, *Interval exchange transformations and measured foliations*, Ann. Math. 115 (1982), 169–200.
[McM03] C. McMullen, *Teichmüller geodesics of infinite complexity*, Acta Math. 191 (2003), 191–223.
[McM03b] C. McMullen, *Billiards and Teichmüller curves on Hilbert modular surfaces*, J. Amer. Math. Soc. 16 (2003), 857–885.
[MW15] M. Mirzakhani, A. Wright, *The boundary of an affine invariant submanifold*, arXiv:1508.01446.
[Mo06] M. Möller, *Variations of Hodge structures of Teichmüller curves*, J. Amer. Math. Soc. 19 (2006), 327–344.
[MV10] M. Möller, E. Viehweg, *Kobayashi geodesics in $\mathcal{A}_g$*, J. Diff. Geom. 86 (2010), 355–379.
[R08] I. Rivin, *Walks on groups, counting reducible matrices, polynomials, and surface and free group automorphisms*, Duke Math. J. 142 (2008), 353–379.
[V86] W. Veech, *The Teichmüller geodesic flow*, Ann. Math. 124 (1990), 441–530.
[VV02] P. Viana, P. M. Veloso, *Galois theory of reciprocal polynomials*, Amer. Math. Monthly 109 (2002), 466–471.
[W14] A. Wright, *The field of definition of affine invariant submanifolds of the moduli space of abelian differentials*. Geom. Top. 18 (2014), 1323–1341.
[W15] A. Wright, *Cylinder deformations in orbit closures of translation surfaces*, Geom. Top. 19 (2015), 413–438.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BONN
ENDENICHER ALLEE 60
53115 BONN, GERMANY
e-mail: ursula@math.uni-bonn.de